

OP MINICHOP

Types: ONCOLOGY TREATMENT

Synonyms: CHOP, LYMPHOMA, LIMP, CYCLOPHOSPHAMIDE, VINCRISTINE, DOXORUBICIN, PREDNISONE, MINI, MINICHOP

Take-Home Medications		Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1	
Take-Home Medications Prior to Treatment			
predniSONE (DELTASONE) 50 MG tablet			
Dose: 40 mg/m ²	Route: oral	daily	
Dispense: --	Refills: --		
Start: S	End: S+5		
Cycles 1 to 6		Repeat 6 times	Cycle length: 21 days
Day 1		Perform every 1 day x1	
Appointment Requests			
INFUSION APPOINTMENT REQUEST			
Interval: --		Occurrences: --	
Labs			
☒ CBC WITH PLATELET AND DIFFERENTIAL			
Interval: --		Occurrences: --	
☒ COMPREHENSIVE METABOLIC PANEL			
Interval: --		Occurrences: --	
☒ MAGNESIUM LEVEL			
Interval: --		Occurrences: --	
☐ LDH			
Interval: --		Occurrences: --	
☐ URIC ACID LEVEL			
Interval: --		Occurrences: --	
☐ ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED			
Interval: --		Occurrences: --	
Outpatient Electrolyte Replacement Protocol			
TREATMENT CONDITIONS 39			
Interval: --		Occurrences: --	
Comments:		Potassium (Normal range 3.5 to 5.0mEq/L)	
		o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
		o Protocol applies only to same day lab value.	
		o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
		o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
		o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
		o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
		o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
		o Sign electrolyte replacement order as Per protocol: cosign required	
TREATMENT CONDITIONS 40			
Interval: --		Occurrences: --	

Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Provider Communication

ONC PROVIDER COMMUNICATION

Interval: -- Occurrences: --

Comments: Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date).

If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.

Nursing Orders

ONC VERIFY PATIENT HAS PRESCRIPTION

Interval: -- Occurrences: --

Comments: Verify that patient has a prescription for oral Prednisone to be started on Days 1-5. If not, please contact physician.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --

Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions: To keep vein open.

Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 100 mL/hr for 1 dose

Start: S

Pre-Medications

☒ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 0.25 mg Route: intravenous once for 1 dose
Offset: 30 Minutes

☐ **dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB**

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Breakthrough Anti-Emetics

☐ **diphenhydramine (BENADRYL) tablet 12.5-25 mg**

Dose: 12.5-25 mg Route: oral every 4 hours PRN
Start: S

☐ **diphenhydramine (BENADRYL) injection 12-25 mg**

Dose: 12-25 mg Route: intravenous every 4 hours PRN
Start: S

☐ **promethazine (PHENERGAN) tablet 12.5-25 mg**

Dose: 12.5-25 mg Route: oral every 4 hours PRN
Start: S

☐ **promethazine (PHENERGAN) injection 12.5 mg**

Dose: 12.5 mg Route: intravenous every 4 hours PRN
Start: S

☐ **promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB**

Dose: 25 mg Route: intravenous every 4 hours PRN over 30 Minutes
Start: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PROMETHAZINE 25 MG/ML	Medications	25 mg	Main Inredient	No

			INJECTION SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
☐	LORazepam (ATIVAN) tablet 0.5-1 mg						
	Dose: 0.5-1 mg Start: S	Route: oral	every 4 hours PRN				
☐	LORazepam (ATIVAN) injection 0.5-1 mg						
	Dose: 0.5-1 mg Start: S	Route: intravenous	every 4 hours PRN				
☐	haloperidol (HALDOL) tablet 0.5-1 mg						
	Dose: 0.5-1 mg Start: S	Route: oral	every 4 hours PRN				
☐	haloperidol lactate (HALDOL) injection 0.5-1 mg						
	Dose: 0.5-1 mg Start: S	Route: intravenous	every 4 hours PRN				
☐	metoclopramide (REGLAN) tablet 10 mg						
	Dose: 10 mg Start: S	Route: oral	every 4 hours PRN				
☐	metoclopramide (REGLAN) injection 10 mg						
	Dose: 10 mg Start: S	Route: intravenous	every 4 hours PRN				
☐	dexamethasone (DECADRON) tablet 10 mg						
	Dose: 10 mg Start: S	Route: oral	every 12 hours PRN				
☐	dexamethasone (DECADRON) injection 10 mg						
	Dose: 10 mg Start: S	Route: intravenous	every 12 hours PRN				
Nursing Orders							
ONC NURSING COMMUNICATION 36							
Interval: --		Occurrences: --					
Comments:		Administer chemotherapy in listed order unless otherwise indicated.					
Chemotherapy							

cyclophosphamide (CYTOXAN) 400 mg/m2 in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 400 mg/m2

Route: intravenous

once over 60 Minutes for 1 dose

Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CYCLOPHOSPHAMIDE 1 GRAM INTRAVENOUS SOLUTION	Medications	400 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

vinCRISTine (ONCOVIN) 1 mg in sodium chloride 0.9% 50 mL chemo IVPB

Dose: 1 mg

Route: intravenous

once over 15 Minutes for 1 dose

Offset: 1.5 Hours

Instructions:

Protect from light, VESICANT. Flat dose of 1 mg.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	VINCRIStINE 1 MG/ML INTRAVENOUS SOLUTION	Medications	1 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

DOXOrubicin (ADRIAMycin) 25 mg/m2 in sodium chloride 0.9% 50 mL chemo IVPB

Dose: 25 mg/m2

Route: intravenous

once over 15 Minutes for 1 dose

Offset: 1.75 Hours

Instructions:

Protect from light; VESICANT

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOXORUBICIN 50 MG/25 ML INTRAVENOUS SOLUTION	Medications	25 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Post-Medications

☐ **pegfilgrastim (NEULASTA) on-body injection
kit 6 mg**

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of
the arm or abdomen (only use the back of the
arm if caregiver is available to monitor
On-body injection status).