

OP MITOXANTRONE / PREDNISONE

Types: ONCOLOGY TREATMENT

Synonyms: NOVA, MITOXANTRONE, NOVANTRONE, PREDNISONE, MITO, PRED, STEROID, PROSTATE

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Take-Home Medications Prior to Treatment		
☉ predniSONE (DELTASONE) 5 MG tablet		
Dose: 5 mg	Route: oral	2 times daily
Dispense: 42 tablet	Refills: 0	
Start: S	End: S+21	
Cycles 1 to 4	Repeat 4 times	Cycle length: 21 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED		
Interval: --	Occurrences: --	
<input type="checkbox"/> PROSTATE SPECIFIC ANTIGEN		
Interval: --	Occurrences: --	
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
	o Sign electrolyte replacement order as Per protocol: cosign required	
TREATMENT CONDITIONS 40		
Interval: --	Occurrences: --	
Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact	

MD/NP

- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL
Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
Start: S End: S 11:15 AM

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

mitoxantrone (NOVANTRONE) 12 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB
Dose: 12 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	MITOXANTRONE 2 MG/ML CONCENTRATE, IN TRAVENOUS	Medications	12 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --

Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEparin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Post-Medications

**pegfilgrastim (NEULASTA) on-body injection
kit 6 mg**

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S End: S

Instructions:

Apply to intact, nonirritated skin on the back of
the arm or abdomen (only use the back of the
arm if caregiver is available to monitor
On-body injection status).