

OP MVAC (DOSE DENSE)

Types: ONCOLOGY TREATMENT

Synonyms: MVAC, METHOTREXATE, VINBLASTINE , DOXORUBICIN , CISPLATIN , VELBAN, ADRIAMYCIN, PLATINOL, BLADDER

Cycles 1 to 6	Repeat 6 times	Cycle length: 14 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: -- Occurrences: --		
<input type="checkbox"/> ZZ_HIST_NM CARDIAC BLOOD POOL MUGA (AT REST)		
Interval: -- Occurrences: --		
<input type="checkbox"/> ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED		
Interval: -- Occurrences: --		
<input type="checkbox"/> ZZ_HIST_MRI CARDIAC FOR MORPH W		
Interval: -- Occurrences: --		
Provider Communication		
ONC PROVIDER COMMUNICATION		
Interval: -- Occurrences: --		
Comments: Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date).		
If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient		

Electrolyte Replacement"

- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: --

Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL

Route: intravenous

once @ 500 mL/hr for 1 dose

Start: S

Pre-Medications

ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
 Start: S End: S 11:15 AM

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
 Start: S

ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

methotrexate PF 30 mg/m2 in sodium chloride 0.9 % 50 mL IVPB

Dose: 30 mg/m2 Route: intravenous once over 15 Minutes for 1 dose
 Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	METHOTREXATE SODIUM (PF) 25 MG/ML INJECTION SOLUTION	Medications	30 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	50 mL	No	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
 Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
 Start: S

Instructions:
 Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

ZZ_HIST_NM CARDIAC BLOOD POOL MUGA (AT REST)

Interval: -- Occurrences: --

ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED

Interval: -- Occurrences: --

ZZ_HIST_MRI CARDIAC FOR MORPH W

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments:

- Potassium (Normal range 3.5 to 5.0mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments:

- Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL

Route: intravenous

once @ 500 mL/hr for 1 dose

Start: S

Pre-Medications

 palonosetron (ALOXI) injection 0.25 mg

Dose: 0.25 mg

Route: intravenous

once for 1 dose

Start: S

End: S 1:45 PM

 dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB

Dose: 12 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**

DEXAMETHASONE

Medications

12 mg

Main

Yes

4 MG/ML

INJECTION

SOLUTION

SODIUM

Base

50 mL

Yes

Yes

CHLORIDE 0.9 %

INTRAVENOUS

SOLUTION

DEXTROSE 5 % IN

Base

50 mL

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

 aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg

Route: intravenous

once over 30 Minutes for 1 dose

Start: S

End: S

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**

APREPITANT 7.2

Medications

130 mg

Main

Yes

MG/ML

INTRAVENOUS

EMULSION

DEXTROSE 5 % IN

Base

130 mL

Yes

Yes

WATER (D5W) IV

SOLP (EXCEL;

NON-PVC)

SODIUM

Base

130 mL

No

Yes

CHLORIDE 0.9 % IV

SOLP

(EXCEL;NON-PVC)

 netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule

Dose: 1 capsule

Route: oral

once for 1 dose

Start: S

End: S 5:30 PM

Instructions:

Administer approximately 1 hour prior to chemotherapy.

SOLUTION

CISplatin (PLATINOL) 70 mg/m2 in sodium chloride 0.9% 500 mL chemo IVPB

Dose: 70 mg/m2 Route: intravenous once over 2 Hours for 1 dose
Offset: 2.5 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CISPLATIN 1 MG/ML	Medications	70 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION				
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Post-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose
Offset: 4.5 Hours

Instructions:
Following chemotherapy.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Supportive Care

pegfilgrastim (NEULASTA) on-body injection kit 6 mg

Dose: 6 mg Route: subcutaneous once for 1 dose
Start: S End: S

Instructions:
Apply to intact, nonirritated skin on the back of the arm or abdomen (only use the back of the arm if caregiver is available to monitor On-body injection status).