

## OP NAB-PACLITAXEL

Types: ONCOLOGY TREATMENT

Synonyms: ABRAXANE , NAB-PACLITAXEL, NAP, ABE, BRAX, BREAST

Cycles 1 to 4	Repeat 4 times	Cycle length: 21 days
<b>Day 1</b>		Perform every 1 day x1
<b>Appointment Requests</b>		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: -- Occurrences: --		
<b>Labs</b>		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>URINALYSIS, AUTOMATED WITH MICROSCOPY</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>CANCER ANTIGEN 27-29 (CA BR)</b>		
Interval: -- Occurrences: --		
<b>Outpatient Electrolyte Replacement Protocol</b>		
<b>TREATMENT CONDITIONS 39</b>		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
<b>TREATMENT CONDITIONS 40</b>		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or greater, do not give		

- magnesium replacement
  - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
  - o Sign electrolyte replacement order as Per protocol: cosign required

**Nursing Orders**

**TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

**Line Flush**

**sodium chloride 0.9 % flush 20 mL**  
 Dose: 20 mL Route: intravenous PRN  
 Start: S

**Nursing Orders**

**sodium chloride 0.9 % infusion 250 mL**  
 Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S  
 Instructions:  
 To keep vein open.

**Pre-Medications**

**ondansetron (ZOFTRAN) injection 8 mg**

Dose: 8 mg Route: intravenous once for 1 dose  
 Start: S End: S 11:15 AM

**ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose  
 Start: S

**ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB**

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose  
 Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

**Pre-Medications**

**diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose  
 Start: S  
 Instructions:  
 Administer via slow IV push 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9 % 50 mL IVPB**

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose  
 Start: S End: S 11:45 AM  
 Instructions:  
 Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

**diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg      Route: oral      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg      Route: oral      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg      Route: intravenous      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg      Route: oral      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg      Route: oral      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**Supportive Care**

**LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg      Route: intravenous      once PRN  
Start: S

**LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg      Route: oral      once PRN  
Start: S

**Antiemetics**

**promethazine (PHENERGAN) injection 12.5 mg**

Dose: 12.5 mg      Route: injection      once PRN  
Start: S

**Chemotherapy**

**NAB-PAClitaxel (ABRAXANE) chemo infusion  
260 mg/m2 (Treatment Plan)**

Dose: 260 mg/m<sup>2</sup>      Route: intravenous      once over 30 Minutes for 1 dose  
Offset: 30 Minutes

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --      Occurrences: --  
Comments:      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

Post-Medications

**pegfilgrastim (NEULASTA) on-body injection  
kit 6 mg**

Dose: 6 mg      Route: subcutaneous      once for 1 dose

Start: S      End: S

Instructions:

Apply to intact, nonirritated skin on the back of  
the arm or abdomen (only use the back of the  
arm if caregiver is available to monitor  
On-body injection status).