

## OP OBINUTUZUMAB (CYCLE 2-6)

Types: ONCOLOGY TREATMENT

Synonyms: OBINUTUZUMAB , GAZYVA, GAYZVA, CLL, LEUKE

Cycles 2 to 6	Repeat 5 times	Cycle length: 28 days
<b>Day 1</b>		Perform every 1 day x1
<b>Appointment Requests</b>		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: -- Occurrences: --		
<b>Labs</b>		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>LDH</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>URIC ACID LEVEL</b>		
Interval: -- Occurrences: --		
<b>Outpatient Electrolyte Replacement Protocol</b>		
<b>TREATMENT CONDITIONS 39</b>		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
<b>TREATMENT CONDITIONS 40</b>		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or areater. do not give		

magnesium replacement  
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"  
o Sign electrolyte replacement order as Per protocol: cosign required

**Nursing Orders**

**TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

**Line Flush**

**sodium chloride 0.9 % flush 20 mL**  
Dose: 20 mL Route: intravenous PRN  
Start: S

**Provider Communication**

**ONC PROVIDER COMMUNICATION 58**

Interval: -- Occurrences: --  
Comments: Prior to beginning Obinutuzumab infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: Push F2:11554001 drawn on \*\*\*.

**Nursing Orders**

**sodium chloride 0.9 % infusion 250 mL**  
Dose: 250 mL Route: intravenous once @ 50 mL/hr for 1 dose  
Start: S  
Instructions: To keep vein open.

**Pre-Medications**

**ondansetron (ZOFRAN) tablet 8 mg**  
Dose: 8 mg Route: oral once for 1 dose  
Start: S  
Instructions: Administer 30 minutes before Chlorambucil.

**ondansetron (ZOFRAN) injection 8 mg**  
Dose: 8 mg Route: intravenous once for 1 dose  
Start: S  
Instructions: Administer 30 minutes before Chlorambucil.

**Nursing Orders**

**ONC NURSING COMMUNICATION 116**

Interval: -- Occurrences: --  
Comments: Please give patients home med: Chlorambucil (Leukeran) and document on the MAR.

**Pre-Medications**

**acetaminophen (TYLENOL) tablet 650 mg**  
Dose: 650 mg Route: oral once for 1 dose  
Start: S  
Instructions: Give 30 minutes prior to Obinutuzumab.

**Chemotherapy**

**obinutuzumab (GAZYVA) 1,000 mg in sodium chloride 0.9 % 250 mL IVPB**  
Dose: 1,000 mg Route: intravenous once over 4 Hours for 1 dose  
Start: S End: S 6:26 PM

**Instructions:**

Gently invert to mix; do not shake or freeze.

Infusion rate:

Cycle 2-6: If no reaction to previous infusion and the final infusion rate was 100 mg/hour or faster, initiate infusion at 100 mg/hour for 30 minutes; if tolerated, may escalate infusion rate in increments of 100 mg/hour every 30 minutes to a maximum rate of 400 mg/hour.

<b>Ingredients:</b>	<b>Name</b>	<b>Type</b>	<b>Dose</b>	<b>Selected</b>	<b>Adds Vol.</b>
	OBINUTUZUMAB 1,000 MG/40 ML INTRAVENOUS SOLUTION	Medications	1,000 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	210 mL	Yes	Yes

**Hematology & Oncology Hypersensitivity Reaction Standing Order**

**ONC NURSING COMMUNICATION 82**

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or

otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 83**

Interval: -- Occurrences: --  
Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)  
1. Stop the infusion.  
2. Notify the CERT team and treating physician immediately.  
3. Place the patient on continuous monitoring.  
4. Obtain vital signs.  
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.  
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.  
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.  
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.  
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.  
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous PRN  
Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg Route: oral PRN  
Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg Route: intravenous PRN  
Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg Route: intravenous PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg Route: intravenous PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg Route: subcutaneous PRN  
Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: -- Occurrences: --  
Comments: Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Day 2** Perform every 1 day x1

Appointment Request for Neulasta/Neupogen

**INJECTION APPOINTMENT REQUEST 17**

Interval: -- Occurrences: --

Post-Medications

**pegfilgrastim (NEULASTA) injection 6 mg**

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S

**Day 15** Perform every 7 days x1

Appointment Requests

**INFUSION APPOINTMENT REQUEST**

Interval: -- Occurrences: --

Labs

**CBC WITH PLATELET AND DIFFERENTIAL**

Interval: -- Occurrences: --

**COMPREHENSIVE METABOLIC PANEL**

Interval: -- Occurrences: --

**MAGNESIUM LEVEL**

Interval: -- Occurrences: --

**LDH**

Interval: -- Occurrences: --

**URIC ACID LEVEL**

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: -- Occurrences: --

Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**TREATMENT CONDITIONS 40**

Interval: -- Occurrences: --

Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP

- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

**TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Pre-Medications

**ondansetron (ZOFRAN) tablet 8 mg**

Dose: 8 mg                      Route: oral                      once for 1 dose  
 Start: S  
 Instructions:  
 Administer 30 minutes before Chlorambucil.

**ondansetron (ZOFRAN) injection 8 mg**

Dose: 8 mg                      Route: intravenous                      once for 1 dose  
 Start: S  
 Instructions:  
 Administer 30 minutes before Chlorambucil.

Nursing Orders

**ONC NURSING COMMUNICATION 116**

Interval: -- Occurrences: --  
 Comments: Please give patients home med: Chlorambucil (Leukeran) and document on the MAR.