

OP OFATUMUMAB

Types: ONCOLOGY TREATMENT

Synonyms: CHRON, LYMPH, LEUKE, CLL, OFAT, ARZE, ARZ, ARZERRA

Cycle 1	Repeat 1 time	Cycle length: 28 days
Day 1	Perform every 1 day x1	
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: -- Occurrences: --		
<input type="checkbox"/> LDH		
Interval: -- Occurrences: --		
<input type="checkbox"/> URIC ACID LEVEL		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
TREATMENT CONDITIONS 40		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or areater. do not give		

magnesium replacement
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL
Dose: 20 mL Route: intravenous PRN
Start: S

Provider Communication

ONC PROVIDER COMMUNICATION 58
Interval: -- Occurrences: --
Comments: Prior to beginning Ofatumumab infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: Push F2:11554001 drawn on ***.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Nursing Orders

ONC NURSING COMMUNICATION 116
Interval: -- Occurrences: --
Comments: Please give patients home med: Chlorambucil (Leukeran) and document on the MAR.

Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg
Dose: 650 mg Route: oral once for 1 dose
Start: S
Instructions:
Administer 30 minutes prior to ofatumumab.

diphenhydramine (BENADRYL) injection 25 mg
Dose: 25 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer via slow IV push 30 minutes prior to ofatumumab.

hydrocortisone sodium succinate (Solu-CORTEF) injection 400 mg
Dose: 400 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer via slow IV push 30 minutes prior to ofatumumab.

Chemotherapy

ofatumumab (ARZERRA) 300 mg in sodium chloride 0.9 % 1,000 mL IVPB
Dose: 300 mg Route: intravenous once for 1 dose
Offset: 30 Minutes
Instructions:
Infuse per manufacturer recommended infusion instructions. Protect from light.

Do not administer until at least 30 min after pre-medications.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	OFATUMUMAB 100 MG/5 ML INTRAVENOUS SOLUTION	100 Medications	300 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	985 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse,

- loss of consciousness, or incontinence)
1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
 6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
 7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
 8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
 9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
 10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

LDH

Interval: -- Occurrences: --

URIC ACID LEVEL

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments:

- Potassium (Normal range 3.5 to 5.0mEq/L)
 - o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
 - o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
 - o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
 - o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments:

- Magnesium (Normal range 1.6 to 2.6mEq/L)
 - o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
 - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
 - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
 - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

 acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to ofatumumab.

 diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose

Start: S

Instructions:

Administer via slow IV push 30 minutes prior to ofatumumab.

 hydrocortisone sodium succinate (Solu-CORTEF) injection 400 mg

Dose: 400 mg Route: intravenous once for 1 dose

Start: S

Instructions:

Administer via slow IV push 30 minutes prior to ofatumumab.

Chemotherapy

ofatumumab (ARZERRA) 1,000 mg in sodium chloride 0.9 % 1,000 mL IVPBDose: 1,000 mg Route: intravenous once for 1 dose
Offset: 30 Minutes

Instructions:

Infuse per manufacturer recommended infusion instructions. Protect from light.

Do not administer until at least 1 hour after pre-medications.

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
OFATUMUMAB 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	1,000 mg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	950 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.

6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25

mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg

Route: oral

PRN

Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN

Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN

Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN

Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --

Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Cycles 2 to 4

Repeat 3 times

Cycle length: 28 days

Day 1

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

LDH

Interval: -- Occurrences: --

URIC ACID LEVEL

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

- Interval: -- Occurrences: --
- Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
 - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
 - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
 - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL
 Dose: 20 mL Route: intravenous PRN
 Start: S

Provider Communication

ONC PROVIDER COMMUNICATION 58
 Interval: -- Occurrences: --
 Comments: Prior to beginning Ofatumumab infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: Push F2:11554001 drawn on ***.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
 Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Nursing Orders

ONC NURSING COMMUNICATION 116
 Interval: -- Occurrences: --
 Comments: Please give patients home med: Chlorambucil (Leukeran) and document on the MAR.

Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg
 Dose: 650 mg Route: oral once for 1 dose

Start: S
Instructions:
Administer 30 minutes prior to ofatumumab.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer via slow IV push 30 minutes prior to ofatumumab.

hydrocortisone sodium succinate (Solu-CORTEF) injection 400 mg

Dose: 400 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer via slow IV push 30 minutes prior to ofatumumab.

Chemotherapy

ofatumumab (ARZERRA) 1,000 mg in sodium chloride 0.9 % 1,000 mL IVPB

Dose: 1,000 mg Route: intravenous once for 1 dose
Offset: 30 Minutes

Instructions:
Infuse per manufacturer recommended infusion instructions. Protect from light.

Do not administer until at least 1 hour after pre-medications.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	OFATUMUMAB 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	1,000 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	950 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --
Comments:

Occurrences: --
Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --
Comments:

Occurrences: --
Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)
1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT
injection syringe 0.3 mg**

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.