

## OP PEMBROLIZUMAB (EVERY 21 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: KEYT, KEYTRUDA, PEMBROLIZ, PAMBRO, NSCL, LUNG, MELANOMA

Cycles 1 to 4	Repeat 4 times	Cycle length: 21 days
<b>Day 1</b>		Perform every 1 day x1
<b>Appointment Requests</b>		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: -- Occurrences: --		
<b>Labs</b>		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>THYROID STIMULATING HORMONE</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>T3, FREE</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>T4, FREE</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>LDH</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>URIC ACID LEVEL</b>		
Interval: -- Occurrences: --		
<b>Outpatient Electrolyte Replacement Protocol</b>		
<b>TREATMENT CONDITIONS 39</b>		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
<b>TREATMENT CONDITIONS 40</b>		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact		

MD/NP

- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

**TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN  
Start: S

Nursing Orders

**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
Start: S  
Instructions:  
To keep vein open.

Pre-Medications

**diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose  
Start: S  
Instructions:  
Administer via slow IV push 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9 % 50 mL IVPB**

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose  
Start: S End: S 11:45 AM  
Instructions:  
Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

**diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg                      Route: oral                      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg                      Route: oral                      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg                      Route: intravenous                      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg                      Route: oral                      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg                      Route: oral                      once for 1 dose  
Offset: 0 Hours

Instructions:  
Administer 30 minutes prior to chemotherapy.

**Chemotherapy**

**pembrolizumab 2 mg/kg in sodium chloride 0.9 % 100 mL IVPB**

Dose: 2 mg/kg                      Route: intravenous                      once over 30 Minutes for 1 dose  
Start: S                      End: S 2:58 AM

Instructions:  
Administer with low protein binding 0.22 micron in-line filter. Do not shake. Do not mix with other medications.

<b>Ingredients:</b>	<b>Name</b>	<b>Type</b>	<b>Dose</b>	<b>Selected</b>	<b>Adds Vol.</b>
	PEMBROLIZUMAB 25 MG/ML INTRAVENOUS SOLUTION	Medications	2 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes

**Hematology & Oncology Hypersensitivity Reaction Standing Order**

**ONC NURSING COMMUNICATION 82**

Interval: --                      Occurrences: --  
Comments:                      Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)  
1. Stop the infusion.  
2. Place the patient on continuous monitoring.  
3. Obtain vital signs.  
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.  
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.  
6. If less than 30 minutes since the last dose of Diphenhydramine,

administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O<sub>2</sub> saturation of greater than or equal to 92%.

6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.

7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O<sub>2</sub> saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.

6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O<sub>2</sub> saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg

Route: intravenous

PRN

Start: S

#### **fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg

Route: oral

PRN

Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg                      Route: intravenous                      PRN  
Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg                      Route: intravenous                      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg                      Route: intravenous                      PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg                      Route: subcutaneous                      PRN  
Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --                      Occurrences: --  
Comments:                      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL                      Route: intravenous                      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units                      Route: intra-catheter                      once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.