

OP TCH (MAINTENANCE REGIMEN)

Types: ONCOLOGY TREATMENT

Synonyms: TCH, DOCETAXEL, CARBOPLATIN, TRASTUZUMAB, HERCEPTIN, TAXOTERE, PARAPLATIN, TACKS, TRASH, CARB, DOS, BREAST

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Pre-Medications		
<input type="radio"/> dexamethasone (DECADRON) 4 MG tablet Dose: 8 mg Route: oral 2 times daily Dispense: 12 tablet Refills: 5 Start: S End: S+3 Instructions: Start day prior to chemotherapy administration.		
Cycles 2 to 6	Repeat 5 times	Cycle length: 21 days
Day 1		Perform every 1 day x1
Appointment Requests		
<input type="checkbox"/> INFUSION APPOINTMENT REQUEST Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL Interval: -- Occurrences: --		
<input type="checkbox"/> URINALYSIS, AUTOMATED WITH MICROSCOPY Interval: -- Occurrences: --		
<input type="checkbox"/> CANCER ANTIGEN 27-29 (CA BR) Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39 Interval: -- Occurrences: -- Comments: Potassium (Normal range 3.5 to 5.0mEq/L) <ul style="list-style-type: none"> o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP o Protocol applies only to same day lab value. o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement" o Sign electrolyte replacement order as Per protocol: cosign required 		
TREATMENT CONDITIONS 40		

Interval: -- Comments:	Occurrences: -- Magnesium (Normal range 1.6 to 2.6mEq/L) o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP o Protocol applies only to same day lab value. o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement" o Sign electrolyte replacement order as Per protocol: cosign required
---------------------------	--

Nursing Orders

ONC NURSING COMMUNICATION 8

Interval: -- Comments:	Occurrences: -- Verify that patient took DEXAMETHASONE orally prior to chemotherapy. Otherwise, please contact physician to order Dexamethasone IV.
---------------------------	--

Provider Communication

ONC PROVIDER COMMUNICATION

Interval: -- Comments:	Occurrences: -- Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date).
---------------------------	--

If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.

Nursing Orders

TREATMENT CONDITIONS 9

Interval: -- Comments:	Occurrences: -- HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; CrCl LESS than 60
---------------------------	---

Line Flush

sodium chloride 0.9 % flush 20 mL			
Dose: 20 mL	Route: intravenous	PRN	
Start: S			

Nursing Orders

sodium chloride 0.9 % infusion 250 mL			
Dose: 250 mL	Route: intravenous	once @ 30 mL/hr for 1 dose	
Start: S			
Instructions:	To keep vein open.		

Pre-Medications

○	ondansetron (ZOFTRAN) 4 mg/2 mL injection 8 mg		
Dose: 8 mg	Route: intravenous	once for 1 dose	
Start: S	End: S 11:15 AM		

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

ondansetron (ZOFTRAN) 4 mg/2 mL 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Pre-Medications

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Supportive Care

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous once PRN
Start: S

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral once PRN
Start: S

Antiemetics

promethazine (PHENERGAN) injection 12.5 mg

Dose: 12.5 mg Route: injection once PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 36

Interval: -- Occurrences: --
Comments: Administer chemotherapy in listed order unless otherwise indicated.

Chemotherapy

trastuzumab (HERCEPTIN) 6 mg/kg in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 6 mg/kg Route: intravenous once over 30 Minutes for 1 dose

Offset: 30 Minutes

Instructions:
NOT compatible with D5W.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	TRASTUZUMAB 150 MG INTRAVENOUS SOLUTION	Medications	6 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

DOCEtaxel (TAXOTERE) 75 mg/m2 in sodium chloride (NON-PVC) 0.9 % 250 mL chemo IVPB

Dose: 75 mg/m2 Route: intravenous once over 60 Minutes for 1 dose
Offset: 1 Hours

Instructions:
Administer through non-DEHP tubing; Use within 4 hours of preparation; Protect from light.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOCETAXEL 80 MG/4 ML (20 MG/ML) INTRAVENOUS SOLUTION	Medications	75 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	QS Base	250 mL	No	Yes

CARBOplatin (PARAplatin) in sodium chloride 0.9 % 250 mL chemo IVPB

AUC: 6 Use AUC Route: intravenous once over 60 Minutes for 1 dose
Offset: 2 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CARBOPLATIN 10 MG/ML INTRAVENOUS SOLUTION	Medications		Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
 Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
 1. Stop the infusion.
 2. Place the patient on continuous monitoring.
 3. Obtain vital signs.
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.

5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --
 Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: -- Occurrences: --
 Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 ma Route: intravenous PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN

Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN

Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN

Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN

Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --

Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Post-Medications

pegfilgrastim (NEULASTA) on-body injection kit 6 mg

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S End: S

Instructions:

Apply to intact, nonirritated skin on the back of the arm or abdomen (only use the back of the arm if caregiver is available to monitor On-body injection status).