

## OP TRASTUZUMAB / CAPECITABINE (CYCLE 2 AND BEYOND)

Types: ONCOLOGY TREATMENT

Synonyms: TRAS, TRASTUZUMAB, HERCEPTIN, CAPECITABINE, XELODA, HER, CAP, ZELODA, XEL, BREAST

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
<b>Day 1</b>		Perform every 1 day x1
Take-Home Medications Prior to Treatment		
<b>capecitabine (XELODA) 500 mg chemo tablet</b>		
Dose: 1,000 mg/m2	Route: oral	2 times daily
Dispense: --	Refills: --	
Start: S	End: S+14	
<b>Cycles 2 to 5</b>	Repeat 4 times	Cycle length: 21 days
<b>Day 1</b>		Perform every 1 day x1
Appointment Requests		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: --	Occurrences: --	
<input type="checkbox"/> <b>URINALYSIS, AUTOMATED WITH MICROSCOPY</b>		
Interval: --	Occurrences: --	
<input type="checkbox"/> <b>CANCER ANTIGEN 27-29 (CA BR)</b>		
Interval: --	Occurrences: --	
Outpatient Electrolyte Replacement Protocol		
<b>TREATMENT CONDITIONS 39</b>		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
	o Sign electrolyte replacement order as Per protocol: cosign required	
<b>TREATMENT CONDITIONS 40</b>		
Interval: --	Occurrences: --	
Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	

- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

#### Nursing Orders

##### **ONC NURSING COMMUNICATION 15**

Interval: -- Occurrences: --  
 Comments: Verify that the patient has taken appropriate oral chemotherapy medication from home prescription.

#### Provider Communication

##### **ONC PROVIDER COMMUNICATION**

Interval: -- Occurrences: --  
 Comments: Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: \*\*\*% on \*\*\* (date).

If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.

#### Nursing Orders

##### **TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

#### Line Flush

##### **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN  
 Start: S

#### Nursing Orders

##### **sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S  
 Instructions: To keep vein open.

#### Pre-Medications

**diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose  
 Start: S  
 Instructions: Administer via slow IV push 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9 % 50 mL IVPB**

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose

Start: S End: S 11:45 AM

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

**diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg Route: intravenous once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

Supportive Care

**LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg Route: intravenous once PRN  
Start: S

**LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg Route: oral once PRN  
Start: S

Antiemetics

**promethazine (PHENERGAN) injection 12.5 mg**

Dose: 12.5 mg Route: injection once PRN  
Start: S

Chemotherapy

**trastuzumab (HERCEPTIN) 6 mg/kg in sodium chloride 0.9 % 250 mL chemo IVPB**

Dose: 6 mg/kg      Route: intravenous      once over 30 Minutes for 1 dose

Instructions:

NOT compatible with D5W.

**Ingredients:**

<b>Name</b>	<b>Type</b>	<b>Dose</b>	<b>Selected</b>	<b>Adds Vol.</b>
TRASTUZUMAB 150 MG INTRAVENOUS SOLUTION	Medications	6 mg/kg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --      Occurrences: --  
Comments:      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.