

## OP TRASTUZUMAB / OXALIPLATIN / LEUCOVORIN / FLUOROURACIL

*Types:* ONCOLOGY TREATMENT

*Synonyms:* TRASTUZUMAB, HERCEPTIN, HER, TRAS, GE JUNCTION, GASTROESOPHAGEAL, OXALIP, ELOX, LEUCO, LUKE, FLOU, FLUO, 5FU

<b>Cycles 1 to 6</b>	Repeat 6 times	Cycle length: 14 days
<b>Day 1</b>	Perform every 1 day x1	
<b>Appointment Requests</b>		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: -- Occurrences: --		
<b>Labs</b>		
<input checked="" type="checkbox"/> <b>URINALYSIS, AUTOMATED WITH MICROSCOPY</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: -- Occurrences: --		
<input type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: -- Occurrences: --		
<b>Outpatient Electrolyte Replacement Protocol</b>		
<b>TREATMENT CONDITIONS 39</b>		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
<b>TREATMENT CONDITIONS 40</b>		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement		

- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Provider Communication

**ONC PROVIDER COMMUNICATION**

Interval: -- Occurrences: --  
 Comments: Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: \*\*\*% on \*\*\* (date).

If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.

Nursing Orders

**TREATMENT CONDITIONS 4**

Interval: -- Occurrences: --  
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

Line Flush

**dextrose 5% flush syringe 20 mL**

Dose: 20 mL Route: intravenous PRN  
 Start: S  
 Instructions: Administer ONLY for Oxaliplatin.

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN  
 Start: S  
 Instructions: Do NOT administer with Oxaliplatin.

Nursing Orders

**dextrose 5% infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S  
 Instructions: To keep vein open for Oxaliplatin.

**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S  
 Instructions: To keep vein open. Do NOT administer with Oxaliplatin.

Pre-Medications

**diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose  
 Start: S  
 Instructions: Administer via slow IV push 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose

Start: S End: S 11:45 AM

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

**diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) injection 20 mg**

Dose: 20 mg Route: intravenous once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

**acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg Route: oral once for 1 dose  
Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

Supportive Care

**LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg Route: intravenous once PRN  
Start: S

**LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg Route: oral once PRN  
Start: S

Antiemetics

**promethazine (PHENERGAN) injection 12.5 mg**

Dose: 12.5 mg Route: injection once PRN  
Start: S



Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	TRASTUZUMAB 150 MG INTRAVENOUS SOLUTION	Medications	4 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

**OXALIPlatin (ELOXATIN) 85 mg/m2 in dextrose  
5% 500 mL chemo IVPB**

Dose: 85 mg/m2      Route: intravenous      once over 2 Hours for 1 dose  
Offset: 1 Hours

Instructions:

Irritant - avoid extravasation. Flush line with  
D5W before and after oxaliplatin infusion.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	OXALIPLATIN 100 MG/20 ML INTRAVENOUS SOLUTION	Medications	85 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes

**leucovorin 400 mg/m2 in dextrose 5% 100 mL  
chemo IVPB**

Dose: 400 mg/m2      Route: intravenous      once over 2 Hours for 1 dose  
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	LEUCOVORIN CALCIUM 350 MG SOLUTION FOR INJECTION	Medications	400 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	No	Yes

**fluorouracil (ADRUCIL) 2,000 mg/m2 in sodium  
chloride 0.9 % 100 mL chemo infusion -  
AMBULATORY PUMP**

Dose: 2,000 mg/m2      Route: intravenous      once over 46 Hours for 1 dose  
Offset: 3 Hours

Instructions:

Nurses to chart as GIVEN on the MAR.  
Administer via CADD pump.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	FLUOROURACIL 5 GRAM/100 ML INTRAVENOUS SOLUTION	Medications	2,000 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	100 mL	No	Yes
	SODIUM CHLORIDE 0.9 %	QS Base	100 mL	Yes	Yes

INTRAVENOUS  
SOLUTION

Nursing Orders

**ONC NURSING COMMUNICATION 11**

Interval: -- Occurrences: --  
Comments: Exposure to cold may exacerbate oxaliplatin-induced neuropathy (including pharyngolaryngeal dysesthesia). Encourage patient to keep blanket on their chest and/or throat during oxaliplatin infusion. Educate patient to avoid cold drinks/foods, ice chips or exposure to cold water or air for 7 days after oxaliplatin infusion.

**ONC NURSING COMMUNICATION 12**

Interval: -- Occurrences: --  
Comments: Assess and notify provider for persistent neuropathy (Grade 2).

**Day 3**

Perform every 1 day x1

Appointment Requests

**ONC PUMP DISCONNECT APPOINTMENT  
REQUEST**

Interval: -- Occurrences: --

Discharge Nursing Orders

**DISCONNECT CONTINUOUS INFUSION PUMP**

Interval: -- Occurrences: --  
Comments: Disconnect patient from continuous infusion pump.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.