

OP TRASTUZUMAB / PERTUZUMAB / DOCETAXEL (CYCLE 1 ONLY)

Types: ONCOLOGY TREATMENT

Synonyms: TRAS, TRASTUZUMAB, HERCEPTIN, PERTUZUMAB, DOCETAXEL, TAXOTERE, PERJETA, HER, DOCE, TAXO, PER, BREAST

Take-Home Medications		Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1	
Pre-Medications			
☐ dexamethasone (DECADRON) 4 MG tablet			
Dose: 8 mg		Route: oral	2 times daily
Dispense: 12 tablet		Refills: 5	
Start: S		End: S+3	
Instructions:			
Start day prior to chemotherapy administration.			
Cycle 1		Repeat 1 time	Cycle length: 21 days
Day 1		Perform every 1 day x1	
Appointment Requests			
INFUSION APPOINTMENT REQUEST			
Interval: --		Occurrences: --	
Labs			
☒ COMPREHENSIVE METABOLIC PANEL			
Interval: --		Occurrences: --	
☒ CBC WITH PLATELET AND DIFFERENTIAL			
Interval: --		Occurrences: --	
☒ MAGNESIUM LEVEL			
Interval: --		Occurrences: --	
☐ URINALYSIS, AUTOMATED WITH MICROSCOPY			
Interval: --		Occurrences: --	
☐ CANCER ANTIGEN 27-29 (CA BR)			
Interval: --		Occurrences: --	
Outpatient Electrolyte Replacement Protocol			
TREATMENT CONDITIONS 39			
Interval: --		Occurrences: --	
Comments:		Potassium (Normal range 3.5 to 5.0mEq/L)	
		o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
		o Protocol applies only to same day lab value.	
		o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
		o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
		o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
		o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
		o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
		o Sign electrolyte replacement order as Per protocol: cosign required	
TREATMENT CONDITIONS 40			

Interval: --
Comments:

Occurrences: --
Magnesium (Normal range 1.6 to 2.6mEq/L)
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
o Protocol applies only to same day lab value.
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 8

Interval: --
Comments:

Occurrences: --
Verify that patient took DEXAMETHASONE orally prior to chemotherapy. Otherwise, please contact physician to order Dexamethasone IV.

Provider Communication

ONC PROVIDER COMMUNICATION

Interval: --
Comments:

Occurrences: --
Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date).

If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --
Comments:

Occurrences: --
HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL
Start: S

Route: intravenous PRN

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL
Start: S

Route: intravenous once @ 30 mL/hr for 1 dose

Instructions:
To keep vein open.

Pre-Medications

● ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg
Start: S

Route: intravenous once for 1 dose
End: S 11:15 AM

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

☐ **ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB**

Dose: 16 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:00 AM

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

Main

No

Ingredient

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

Pre-Medications

☒ **diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg

Route: intravenous

once for 1 dose

Start: S

Instructions:

Administer via slow IV push 30 minutes prior to chemotherapy.

☐ **diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9 % 50 mL IVPB**

Dose: 50 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:45 AM

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

DIPHENHYDRAMIN
E 50 MG/ML
INJECTION
SOLUTION

Medications

50 mg

Main

No

Ingredient

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

Yes

Yes

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

50 mL

No

Yes

☐ **diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg

Route: intravenous

once for 1 dose

Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

☐ **famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg Route: oral once for 1 dose
Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

☒ **acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg Route: oral once for 1 dose
Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

Supportive Care

☐ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg Route: intravenous once PRN
Start: S

☐ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg Route: oral once PRN
Start: S

Antiemetics

☐ **promethazine (PHENERGAN) injection 12.5 mg**

Dose: 12.5 mg Route: injection once PRN
Start: S

Chemotherapy

trastuzumab (HERCEPTIN) 8 mg/kg in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 8 mg/kg Route: intravenous once over 90 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
NOT compatible with D5W.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	TRASTUZUMAB 150 MG INTRAVENOUS SOLUTION	Medications	8 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

pertuzumab (PERJETA) 840 mg in sodium chloride 0.9 % 250 mL IVPB

Dose: 840 mg Route: intravenous once over 1 Hours for 1 dose
Offset: 2 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PERTUZUMAB 420 MG/14 ML (30 MG/ML) INTRAVENOUS SOLUTION	Medications	840 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	222 mL	Yes	Yes

DOCEtaxel (TAXOTERE) 75 mg/m2 in sodium

chloride (NON-PVC) 0.9 % 250 mL chemo IVPB

Dose: 75 mg/m2

Route: intravenous

once over 60 Minutes for 1 dose

Offset: 3 Hours

Instructions:

Administer through non-DEHP tubing; Use within 4 hours of preparation; Protect from light.

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**DOCETAXEL 80
MG/4 ML (20
MG/ML)

Medications

75 mg/m2

Main

Yes

Ingredient

INTRAVENOUS
SOLUTIONSODIUM
CHLORIDE 0.9 % IV

QS Base

250 mL

Yes

Yes

SOLP
(EXCEL;NON-PVC)DEXTROSE 5 % IN
WATER (D5W) IV

QS Base

250 mL

No

Yes

SOLP (EXCEL;
NON-PVC)

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Post-Medications

☐ **pegfilgrastim (NEULASTA) on-body injection kit 6 mg**

Dose: 6 mg

Route: subcutaneous

once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of the arm or abdomen (only use the back of the arm if caregiver is available to monitor On-body injection status).