

OP TRASTUZUMAB / PERTUZUMAB / DOCETAXEL (CYCLE 2 AND BEYOND)

Types: ONCOLOGY TREATMENT

Synonyms: TRAS, TRASTUZUMAB, HERCEPTIN, PERTUZUMAB, DOCETAXEL, TAXOTERE, PERJETA, HER, PER, TAXO, TAX, HUR, HURCEPTIN, BREAST

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Pre-Medications		
<input type="radio"/> dexamethasone (DECADRON) 4 MG tablet		
Dose: 8 mg	Route: oral	2 times daily
Dispense: 12 tablet	Refills: 5	
Start: S	End: S+3	
Instructions: Start day prior to chemotherapy administration.		
Cycles 2 to 5	Repeat 4 times	Cycle length: 21 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
<input type="checkbox"/> URINALYSIS, AUTOMATED WITH MICROSCOPY		
Interval: --	Occurrences: --	
<input type="checkbox"/> CANCER ANTIGEN 27-29 (CA BR)		
Interval: --	Occurrences: --	
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
	o Sign electrolyte replacement order as Per protocol: cosign required	

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --
 Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 8

Interval: -- Occurrences: --
 Comments: Verify that patient took DEXAMETHASONE orally prior to chemotherapy. Otherwise, please contact physician to order Dexamethasone IV.

Provider Communication

ONC PROVIDER COMMUNICATION

Interval: -- Occurrences: --
 Comments: Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date).

If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL
 Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
 Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Pre-Medications

ondansetron (ZOFRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
 Start: S End: S 11:15 AM

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
 Start: S

ondansetron (ZOFRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Pre-Medications

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
 Start: S
 Instructions:
 Administer via slow IV push 30 minutes prior to chemotherapy.

diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9% 50 mL IVPB

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:45 AM
 Instructions:
 Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

diphenhydrAMINE (BENADRYL) tablet 25 mg

Dose: 25 mg Route: oral once for 1 dose
 Offset: 0 Hours
 Instructions:
 Administer 30 minutes prior to chemotherapy.

diphenhydrAMINE (BENADRYL) tablet 50 mg

Dose: 50 mg Route: oral once for 1 dose
Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

famotidine (PEPCID) injection 20 mg

Dose: 20 mg Route: intravenous once for 1 dose
Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

famotidine (PEPCID) tablet 20 mg

Dose: 20 mg Route: oral once for 1 dose
Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose
Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

Supportive Care

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous once PRN
Start: S

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral once PRN
Start: S

Antiemetics

promethazine (PHENERGAN) injection 12.5 mg

Dose: 12.5 mg Route: injection once PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 36

Interval: -- Occurrences: --
Comments: Administer chemotherapy in listed order unless otherwise indicated.

Chemotherapy

trastuzumab (HERCEPTIN) 6 mg/kg in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 6 mg/kg Route: intravenous once over 30 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
NOT compatible with D5W.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	TRASTUZUMAB 150 MG INTRAVENOUS SOLUTION	Medications	6 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

pertuzumab (PERJETA) 420 mg in sodium chloride 0.9 % 250 mL IVPB

Dose: 420 mg Route: intravenous once over 1 Hours for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PERTUZUMAB 420 MG/14 ML (30 MG/ML) INTRAVENOUS SOLUTION	Medications	420 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	236 mL	Yes	Yes

DOCEtaxel (TAXOTERE) 75 mg/m2 in sodium chloride (NON-PVC) 0.9 % 250 mL chemo IVPB

Dose: 75 mg/m2 Route: intravenous once over 60 Minutes for 1 dose
Offset: 2 Hours

Instructions:
Administer through non-DEHP tubing; Use within 4 hours of preparation; Protect from light.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOCETAXEL 80 MG/4 ML (20 MG/ML) INTRAVENOUS SOLUTION	Medications	75 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	QS Base	250 mL	No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN


Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Post-Medications

pegfilgrastim (NEULASTA) on-body injection kit 6 mg

Dose: 6 mg Route: subcutaneous once for 1 dose
Start: S End: S

Instructions:



Apply to intact, nonirritated skin on the back of the arm or abdomen (only use the back of the arm if caregiver is available to monitor On-body injection status).