

OP TRASTUZUMAB / PERTUZUMAB / PACLITAXEL (CYCLE 2 AND BEYOND)

Types: ONCOLOGY TREATMENT

Synonyms: TRAS, TRASTUZUMAB, HERCEPTIN, PERTUZUMAB, PACLITAXEL, TAXOL, PERJETA, HER, PER, TAX, BREAST , THP

Cycles 2 to 4		Repeat 3 times	Cycle length: 21 days
Day 1	Appointment Requests		Perform every 1 day x1
	INFUSION APPOINTMENT REQUEST		
	Interval: --	Occurrences: --	
	Labs		
	☒ COMPREHENSIVE METABOLIC PANEL		
	Interval: --	Occurrences: --	
	☒ CBC WITH PLATELET AND DIFFERENTIAL		
	Interval: --	Occurrences: --	
	☒ MAGNESIUM LEVEL		
	Interval: --	Occurrences: --	
Outpatient	☐ URINALYSIS, AUTOMATED WITH MICROSCOPY		
	Interval: --	Occurrences: --	
	☐ CANCER ANTIGEN 27-29 (CA BR)		
	Interval: --	Occurrences: --	
	Electrolyte Replacement Protocol		
	TREATMENT CONDITIONS 39		
	Interval: --	Occurrences: --	
	Comments:	Potassium (Normal range 3.5 to 5.0mEq/L) - o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP - o Protocol applies only to same day lab value. - o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP - o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO - o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO - o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement" - o Sign electrolyte replacement order as Per protocol: cosign required	
	TREATMENT CONDITIONS 40		
	Interval: --	Occurrences: --	
	Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L) - o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP - o Protocol applies only to same day lab value. - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium	

- sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Pre-Medications

☒ ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
 Start: S End: S 11:15 AM

☐ ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
 Start: S

☐ ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Pre-Medications

☒ diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
 Start: S
 Instructions:
 Administer via slow IV push 30 minutes prior to chemotherapy.

☐ diphenhydramine (BENADRYL) 50 mg in sodium chloride 0.9% 50 mL IVPB

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:45 AM

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:**Name****Type****Dose****Selected**
Main
Ingredient**Adds Vol.**DIPHENHYDRAMIN
E 50 MG/ML
INJECTION
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Medications

50 mg

No

Base

50 mL

Yes

Yes

Base

50 mL

No

Yes

☐ **diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☒ **famotidine (PEPCID) injection 20 mg**

Dose: 20 mg

Route: intravenous

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

Supportive Care☐ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg

Route: intravenous

once PRN

Start: S

☐ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg

Route: oral

once PRN

Start: S

Nursing Orders**ONC NURSING COMMUNICATION 36**

Interval: --

Occurrences: --

Comments:

Administer chemotherapy in listed order unless otherwise indicated.

Chemotherapy

trastuzumab (HERCEPTIN) 6 mg/kg in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 6 mg/kg

Route: intravenous

once over 30 Minutes for 1 dose

Offset: 30 Minutes

Instructions:

NOT compatible with D5W.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

TRASTUZUMAB
150 MG
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Medications

6 mg/kg

Main

Yes

Ingredient

QS Base

250 mL

Yes

Yes

pertuzumab (PERJETA) 420 mg in sodium chloride 0.9 % 250 mL IVPB

Dose: 420 mg

Route: intravenous

once over 1 Hours for 1 dose

Offset: 1 Hours

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

PERTUZUMAB 420
MG/14 ML (30
MG/ML)
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Medications

420 mg

Main

Yes

Ingredient

QS Base

236 mL

Yes

Yes

PAClitaxel (TAXOL) 80 mg/m2 in sodium chloride (NON-PVC) 0.9 % 250 mL chemo IVPB

Dose: 80 mg/m2

Route: intravenous

once over 1 Hours for 1 dose

Offset: 2 Hours

Instructions:

Administer through a 0.22 micron filter and non-PVC tubing set.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

PACLITAXEL 6
MG/ML
CONCENTRATE, IN
TRAVENOUS
SODIUM
CHLORIDE 0.9 % IV
SOLP
(EXCEL; NON-PVC)
DEXTROSE 5 % IN
WATER (D5W) IV
SOLP (EXCEL;
NON-PVC)

Medications

80 mg/m2

Main

Yes

Ingredient

QS Base

250 mL

Yes

Yes

QS Base

No

Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 8

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: -- Occurrences: --

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: -- Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: -- Occurrences: --

☐ **URINALYSIS, AUTOMATED WITH MICROSCOPY**

Interval: -- Occurrences: --

☐ **CANCER ANTIGEN 27-29 (CA BR)**

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments:

Potassium (Normal range 3.5 to 5.0mEq/L)
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
o Protocol applies only to same day lab value.
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
o Protocol applies only to same day lab value.
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement

- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

☒ ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg

Route: intravenous

once for 1 dose

Start: S

End: S 11:15 AM

☐ ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

☐ ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:00 AM

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON

Medications

16 mg

Main

No

HCL (PF) 4 MG/2

Ingredient

ML INJECTION

SOLUTION

DEXTROSE 5 % IN Base

50 mL

Always

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

Pre-Medications

☒ diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

once for 1 dose

Start: S

Instructions:

Administer via slow IV push 30 minutes prior to chemotherapy.

☐ diphenhydramine (BENADRYL) 50 mg in sodium chloride 0.9% 50 mL IVPB

Dose: 50 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:45 AM

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

			DIPHENHYDRAMIN Medications		50 mg	Main	No
			E 50 MG/ML INJECTION SOLUTION			Ingredient	
			SODIUM CHLORIDE 0.9 %	Base	50 mL	Yes	Yes
			INTRAVENOUS SOLUTION				
			DEXTROSE 5 % IN WATER (D5W)	Base	50 mL	No	Yes
			INTRAVENOUS SOLUTION				
			☐ diphenhydrAMINE (BENADRYL) tablet 25 mg				
			Dose: 25 mg	Route: oral	once for 1 dose Offset: 0 Hours		
			Instructions: Administer 30 minutes prior to chemotherapy.				
			☐ diphenhydrAMINE (BENADRYL) tablet 50 mg				
			Dose: 50 mg	Route: oral	once for 1 dose Offset: 0 Hours		
			Instructions: Administer 30 minutes prior to chemotherapy.				
			☒ famotidine (PEPCID) injection 20 mg				
			Dose: 20 mg	Route: intravenous	once for 1 dose Offset: 0 Hours		
			Instructions: Administer 30 minutes prior to chemotherapy.				
			☐ famotidine (PEPCID) tablet 20 mg				
			Dose: 20 mg	Route: oral	once for 1 dose Offset: 0 Hours		
			Instructions: Administer 30 minutes prior to chemotherapy.				
			☐ acetaminophen (TYLENOL) tablet 650 mg				
			Dose: 650 mg	Route: oral	once for 1 dose Offset: 0 Hours		
			Instructions: Administer 30 minutes prior to chemotherapy.				
Supportive Care							
☐ LORAZepam (ATIVAN) injection 1 mg							
Dose: 1 mg	Route: intravenous	once PRN					
Start: S							
☐ LORAZepam (ATIVAN) tablet 1 mg							
Dose: 1 mg	Route: oral	once PRN					
Start: S							
Chemotherapy							
PACLitaxel (TAXOL) 80 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB							
Dose: 80 mg/m2	Route: intravenous	once over 1 Hours for 1 dose Offset: 30 Minutes					
Instructions: Administer through a 0.22 micron filter and non-PVC tubing set.							
Ingredients:		Name	Type	Dose	Selected	Adds Vol.	

PACLITAXEL 6 MG/ML CONCENTRATE, IN TRAVENOUS SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC) DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Medications	80 mg/m2	Main Ingredient	Yes
	QS Base		Yes	Yes
	QS Base		No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

☒ sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

☒ HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 15

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

☒ COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

☒ CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

☒ MAGNESIUM LEVEL

Interval: -- Occurrences: --

☐ URINALYSIS, AUTOMATED WITH MICROSCOPY

Interval: -- Occurrences: --

☐ CANCER ANTIGEN 27-29 (CA BR)

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
o Protocol applies only to same day lab value.
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO

- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: --

Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

● ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg

Route: intravenous

once for 1 dose

Start: S

End: S 11:15 AM

○ ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

○ ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:00 AM

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON

Medications

16 mg

Main

No

HCL (PF) 4 MG/2 ML INJECTION SOLUTION DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
--	------	-------	--------	-----

Ingredient

Pre-Medications

☒ **diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer via slow IV push 30 minutes prior to chemotherapy.

☐ **diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:45 AM
Instructions:
Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

☐ **diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

☐ **diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

☒ **famotidine (PEPCID) injection 20 mg**

Dose: 20 mg Route: intravenous once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

☐ **famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

☐ **acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg Route: oral once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

Supportive Care

☐ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg

Route: intravenous

once PRN

Start: S

☐ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg

Route: oral

once PRN

Start: S

Chemotherapy

PACLitaxel (TAXOL) 80 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 80 mg/m2

Route: intravenous

once over 1 Hours for 1 dose

Offset: 30 Minutes

Instructions:

Administer through a 0.22 micron filter and non-PVC tubing set.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

PACLITAXEL 6
MG/ML
CONCENTRATE, IN
TRAVENOUS
SODIUM
CHLORIDE 0.9 % IV
SOLP
(EXCEL; NON-PVC)
DEXTROSE 5 % IN
WATER (D5W) IV
SOLP (EXCEL;
NON-PVC)

Medications

80 mg/m2

Main
Ingredient

Yes

QS Base

Yes

Yes

QS Base

No

Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.