Protocol Index

OP XELIRI (EVERY 21 DAYS)

Types: ONCOLOGY TREATMENT
Synonyms: XELIRI, CELERY, CAPECITABINE , IRINOTECAN , XELODA, CAMPTOSAR, ZELODA, IRENE, COLORECTAL

Take-Home Medications Repeat 1 time Cycle length: 1 day

Day 1 Perform every 1 day x1

Take-Home Medications Prior to Treatment

capecitabine (XELODA) 500 mg chemo tablet
Dose: 1,000 mg/m2 Route: oral 2 times daily
Dispense: -- Refills: --
Start: S End: S+14

Cycles 1 to 4 Repeat 4 times Cycle length: 21 days

Day 1 Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST
Interval: -- Occurrences: --

Labs

☑ **COMPREHENSIVE METABOLIC PANEL**
Interval: -- Occurrences: --

☑ **CBC WITH PLATELET AND DIFFERENTIAL**
Interval: -- Occurrences: --

☑ **MAGNESIUM LEVEL**
Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39
Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)
- Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- Protocol applies only to same day lab value.
- Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40
Interval: -- Occurrences: --
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- Protocol applies only to same day lab value.
- Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium

sulfate IV
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Nursing Orders

ONC NURSING COMMUNICATION 15

Interval: -- Occurrences: --
Comments: Verify that the patient has taken appropriate oral chemotherapy medication from home prescription.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

☑ **ondansetron (ZOFRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:30 AM

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

☐ **ondansetron (ZOFRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose
Start: S End: S 11:30 AM

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg Route: oral once for 1 dose

Start: S

☐ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 250 mcg | Route: intravenous | once for 1 dose
Start: S | End: S 3:00 PM
Instructions:
For OUTPATIENT use only.

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg | Route: intravenous | once over 30 Minutes for 1 dose
Start: S | End: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

atropine injection 0.25 mg

Dose: 0.25 mg | Route: intravenous | once PRN
Start: S

Supportive Care

○ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg | Route: intravenous | once PRN
Start: S

○ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg | Route: oral | once PRN
Start: S

Chemotherapy

irinotecan (CAMPTOSAR) 250 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 250 mg/m2 | Route: intravenous | once over 90 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
Protect from light

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
IRINOTECAN 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	250 mg/m2	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

☑ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☑ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.