

OP XELIRI / PANITUMUMAB (EVERY 7 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: FOLFOXIRI, FLUOROURACIL , LEUCOVORIN, OXALIPLATIN , IIRINOTECAN , AGE, CAMPTOSAR, ADRUCIL, ELOXATIN, FLOW, IRENE, PANIT, VECTIB

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Take-Home Medications Prior to Treatment		
capecitabine (XELODA) 500 mg chemo tablet		
Dose: 1,000 mg/m ²	Route: oral	2 times daily
Dispense: --	Refills: --	
Start: S	End: S+14	
Cycles 1 to 6	Repeat 6 times	Cycle length: 14 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
	o Sign electrolyte replacement order as Per protocol: cosign required	
TREATMENT CONDITIONS 40		
Interval: --	Occurrences: --	
Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP	
	o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV	
	o Serum Magnesium 1.3 to 1.5mEq/L. give 1 gram magnesium	

- sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Nursing Orders

ONC NURSING COMMUNICATION 15

Interval: -- Occurrences: --
 Comments: Verify that the patient has taken appropriate oral chemotherapy medication from home prescription.

Provider Communication

ONC PROVIDER COMMUNICATION 2

Interval: -- Occurrences: --
 Comments: Tumor KRAS gene status should be determined prior to initiation of therapy. KRAS type: Please Push F2:115540219.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone

- (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:30 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S End: S 11:30 AM

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

atropine injection 0.25 mg

Dose: 0.25 mg Route: intravenous once PRN
Start: S

Supportive Care

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous once PRN
Start: S

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral once PRN
Start: S

Chemotherapy

panitumumab (VECTIBIX) 6 mg/kg in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 6 mg/kg Route: intravenous once over 60 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
USE 0.2 OR 0.22 MICRON INLINE FILTER.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PANITUMUMAB 100 MG/5 ML (20 MG/ML) INTRAVENOUS SOLUTION	Medications	6 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

irinotecan (CAMPTOSAR) 250 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 250 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 1.5 Hours

Instructions:

Protect from light
Ingredients:

Name	Type	Dose	Selected	Adds Vol.
IRINOTECAN 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	250 mg/m2	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 14

Interval: -- Occurrences: --
Comments: Contact Provider if drug-induced acneiform rash develops and covers more than 25 per cent of the body.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.