

OP XELOX / PANITUMUMAB

Types: ONCOLOGY TREATMENT

Synonyms: COLORECTAL, GI, GASTRO, PANIT, VECTI, CAPE, XELOX, XELO, OXA, ELOX

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Take-Home Medications Prior to Treatment		
capecitabine (XELODA) 500 mg chemo tablet		
Dose: 850 mg/m2	Route: oral	2 times daily
Dispense: --	Refills: --	
Start: S	End: S+14	
Cycles 1 to 3	Repeat 3 times	Cycle length: 28 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
	o Sign electrolyte replacement order as Per protocol: cosign required	
TREATMENT CONDITIONS 40		
Interval: --	Occurrences: --	
Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP	
	o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV	
	o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV	

- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Nursing Orders

ONC NURSING COMMUNICATION 15

Interval: -- Occurrences: --
 Comments: Verify that the patient has taken appropriate oral chemotherapy medication from home prescription.

Line Flush

dextrose 5% flush syringe 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S
 Instructions: Administer ONLY for Oxaliplatin.

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S
 Instructions: Do NOT administer with Oxaliplatin.

Nursing Orders

dextrose 5% infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open for Oxaliplatin.

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open. Do NOT administer with Oxaliplatin.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone

- (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:30 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
	SODIUM CHLORIDE 0.9 %	Base	50 mL	Always	Yes

INTRAVENOUS SOLUTION
 DEXTROSE 5 % IN Base No Yes
 WATER (D5W)
 INTRAVENOUS SOLUTION

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
 Start: S End: S 11:30 AM

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
 Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Supportive Care

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous once PRN
 Start: S

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral once PRN
 Start: S

Chemotherapy

panitumumab (VECTIBIX) 6 mg/kg in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 6 mg/kg Route: intravenous once over 60 Minutes for 1 dose
 Offset: 30 Minutes

Instructions:
 USE 0.2 OR 0.22 MICRON INLINE FILTER.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PANITUMUMAB 100 MG/5 ML (20 MG/ML)	Medications	6 mg/kg	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

OXALiPlatin (ELOXATIN) 130 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 130 ma/m2 Route: intravenous once over 120 Minutes for 1 dose

Offset: 1.5 Hours

Instructions:

Irritant - avoid extravasation. Flush line with D5W before and after oxaliplatin infusion.

Ingredients:

Name	Type	Dose	Selected	Adds	Vol.
OXALIPLATIN 100 MG/20 ML INTRAVENOUS SOLUTION	Medications	130 mg/m2	Main Ingredient	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic

compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 11

Interval: -- Occurrences: --
Comments: Exposure to cold may exacerbate oxaliplatin-induced neuropathy (including pharyngolaryngeal dysesthesia). Encourage patient to keep blanket on their chest and/or throat during oxaliplatin infusion. Educate patient to avoid cold drinks/foods, ice chips or exposure to cold water or air for 7 days after oxaliplatin infusion.

ONC NURSING COMMUNICATION 12

Interval: -- Occurrences: --
Comments: Assess and notify provider for persistent neuropathy (Grade 2).

Nursing Orders

ONC NURSING COMMUNICATION 14

Interval: -- Occurrences: --
Comments: Contact Provider if drug-induced acneiform rash develops and covers

more than 25 per cent of the body.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 15

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments:

Potassium (Normal range 3.5 to 5.0mEq/L)

o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP

o Protocol applies only to same day lab value.

o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP

o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO

o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO

o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement

o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"

o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)

o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP

o Protocol applies only to same day lab value.

o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP

- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Nursing Orders

ONC NURSING COMMUNICATION 15

Interval: -- Occurrences: --
 Comments: Verify that the patient has taken appropriate oral chemotherapy medication from home prescription.

Line Flush

dextrose 5% flush syringe 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S
 Instructions:
 Administer ONLY for Oxaliplatin.

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S
 Instructions:
 Do NOT administer with Oxaliplatin.

Nursing Orders

dextrose 5% infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open for Oxaliplatin.

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open. Do NOT administer with Oxaliplatin.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:30 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE	Medications	12 mg	Yes	No
	4 MG/ML				

INJECTION SOLUTION
SODIUM CHLORIDE 0.9 %
INTRAVENOUS SOLUTION
DEXTROSE 5 % IN WATER (D5W)
INTRAVENOUS SOLUTION

Base	50 mL	Always	Yes
Base		No	Yes

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S End: S 11:30 AM

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Supportive Care

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous once PRN
Start: S

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral once PRN
Start: S

Chemotherapy

panitumumab (VECTIBIX) 6 mg/kg in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 6 mg/kg Route: intravenous once over 60 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
USE 0.2 OR 0.22 MICRON INLINE FILTER.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PANITUMUMAB 100 MG/5 ML (20 MG/ML)	Medications	6 mg/kg	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS	QS Base	100 mL	Yes	Yes

SOLUTION

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --
Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: -- Occurrences: --
Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new

bag and new intravenous tubing.
 8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
 9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
 10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
 Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
 Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
 Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
 Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
 Start: S

Nursing Orders

ONC NURSING COMMUNICATION 11

Interval: -- Occurrences: --
 Comments: Exposure to cold may exacerbate oxaliplatin-induced neuropathy (including pharyngolaryngeal dysesthesia). Encourage patient to keep blanket on their chest and/or throat during oxaliplatin infusion. Educate patient to avoid cold drinks/foods, ice chips or exposure to cold water or air for 7 days after oxaliplatin infusion.

ONC NURSING COMMUNICATION 12

Interval: -- Occurrences: --
 Comments: Assess and notify provider for persistent neuropathy (Grade 2).

Nursing Orders

ONC NURSING COMMUNICATION 14

Interval: -- Occurrences: --
 Comments: Contact Provider if drug-induced acneiform rash develops and covers more than 25 per cent of the body.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
 Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.