

**Nursing Orders**

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every 4 weeks	S	Until discontin'td
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Vital Signs PRE & POST Infusion.	Every 4 weeks	S	Until discontin'td
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every 4 weeks	S	Until discontin'td
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every 4 weeks	S	Until discontin'td
<input type="checkbox"/> Discontinue IV Details	Every 4 weeks	S	Until discontin'td

**Medications**

	Interval	Defer Until	Duration
<input type="checkbox"/> belatacept (NULOJIX) 5 mg/kg in sodium chloride 0.9 % 100 mL IVPB 5 mg/kg, intravenous, for 30 Minutes, once, Starting S, For 1 Doses Belatacept is only for use in EBV Seropositive Patients.	Every 4 weeks	S	Until discontin'td