

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> Urinalysis, automated with microscopy STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> Sedimentation rate STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> C-reactive protein STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> C3 complement component STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> C4 complement component STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discount'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Details	Every visit	S	Until discount'd
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for any signs of infusion reaction. For MINOR INFUSION REACTIONS (fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea) stop or slow infusion rate by 50%. If symptoms do not resolve within 5-10 minutes give Benadryl as ordered. Notify MD if symptoms do not resolve within 5-10 minutes. With MD approval, give Solu-medrol as ordered. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor patient closely until completion.	Every visit	S	Until discount'd
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of SEVERE INFUSION REACTIONS: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension. For severe infusion reaction take the following steps; STOP Infusion and start Normal Saline solution, Notify physician and administer Benadryl as ordered, Monitor vital signs, Administer Epinephrine as ordered. Consult MD for further orders.	Every visit	S	Until discount'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discount'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discount'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discount'd

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydramine (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discount'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 500 mg 500 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discount'd
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discount'd

Medications

Medications (continued)

	Interval	Defer Until	Duration
<input type="checkbox"/> belimumab (BENLYSTA) IVPB 10 mg/kg 10 mg/kg, intravenous, once, Starting S, For 1 Doses Administer on Weeks 0, 2, and 4.	Every 2 weeks	S	3 treatments
<input type="checkbox"/> belimumab (BENLYSTA) IVPB 10 mg/kg 10 mg/kg, intravenous, once, Starting S, For 1 Doses Administer on Week 8 and every 4 weeks thereafter.	Every 4 weeks	S+55	Until discontinuation

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours PRN, mild pain (score 1-3), headaches, fever, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 25 mg 25 mg, intravenous, PRN, itching, for mild to moderate infusion reaction, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, For Severe Infusion Reaction, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> epinephrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> methylprednisolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, PRN, for infusion reaction, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discontinuation