BLANK THERAPY PLAN

Nursing Orders

		Interval	Defer Until	Duration
	Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
	Vital signs - T/P/R/BP Per Unit Protocol Details	Every visit	S	Until discont'd
	sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'c
	sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'o
	Discontinue IV Details	Every visit	S	Until discont'c
ersen	sitivity	Intonyal	Defer Until	Duration
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	Interval PRN	S	Until discont'o
	diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S	PRN	S	Until discont'o
	diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S	PRN S, For 1 Doses	S	Until discont'o
	diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting	PRN S, For 1 Doses	S	Until discont'o
	epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'o
	hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Dose	PRN	S	Until discont'o
	promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'o
	sodium chloride 0.9% infusion 250 mL	PRN	S	Until discont'o

250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses