

Labs

	Interval	Defer Until	Duration
<input checked="" type="checkbox"/> Type and screen Routine, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd

Nursing Pre-Transfusion Orders

	Interval	Defer Until	Duration
<input checked="" type="checkbox"/> Consent obtained The Benefits vs Risks of Blood and Blood product transfusion have been discussed with this patient. Has consent been obtained? Yes	Every visit	S	Until discont'd
<input checked="" type="checkbox"/> Maintain IV Access Details	Every visit	S	Until discont'd
<input checked="" type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Monitor Vital Signs before transfusion starts, 15 minutes after transfusion starts, and at completion of transfusion. Notify physician of any abnormal vital signs.	Every visit	S	Until discont'd
<input checked="" type="checkbox"/> Transfusion Reaction Evaluation Transfusion Reaction Evaluation: In the event of a transfusion reaction do the following: 1) Immediately stop the transfusion 2) Start Normal Saline Intravenous infusion 3) Notify ordering physician and Transfusion Medicine Services (blood bank) 4) Document the suspected transfusion reaction (refer to HMH Policy and Procedure NU1303) 5) Obtain post-transfusion type and screen.	Every visit	S	Until discont'd

Transfusion Product

	Interval	Defer Until	Duration
<input checked="" type="checkbox"/> Prepare platelet pheresis Routine Transfusion Indications: Transfusion date:	Every visit	S	Until discont'd
<input checked="" type="checkbox"/> Prepare RBC Routine Transfusion Indications: Transfusion date:	Every visit	S	Until discont'd

Nursing Transfusion Orders

	Interval	Defer Until	Duration
<input checked="" type="checkbox"/> Transfuse platelets Routine Transfusion duration per unit (hrs):	Every visit	S	Until discont'd
<input checked="" type="checkbox"/> Transfuse RBC Routine Transfusion duration per unit (hrs):	Every visit	S	Until discont'd

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discont'd
<input type="checkbox"/> famotidine (PEPCID) tablet 20 mg 20 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discont'd
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg	Every visit	S	Until discont'd

Pre-Medications (continued)

	Interval	Defer Until	Duration
20 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.			
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg 180 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discontin't'd
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discontin't'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 10 mg 10 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discontin't'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to blood transfusion.	Every visit	S	Until discontin't'd

IV Fluids

	Interval	Defer Until	Duration
<input checked="" type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, once, Starting S, For 1 Doses Administer with blood to keep vein open.	Every visit	S	Until discontin't'd
<input type="checkbox"/> sodium chloride 0.9% infusion 500 mL 500 mL, intravenous, at 30 mL/hr, once, Starting S, For 1 Doses Administer with blood to keep vein open.	Every visit	S	Until discontin't'd

Post-Medications

	Interval	Defer Until	Duration
<input checked="" type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S IV flush upon transfusion completion.	Every visit	S	Until discontin't'd
<input type="checkbox"/> furosemide (LASIX) injection 20 mg 20 mg, intravenous, once, For 1 Doses Inject each 20 mg of furosemide slowly IV over 1 to 2 minutes. Administer after first unit of blood.	Every visit	S	Until discontin't'd
<input type="checkbox"/> furosemide (LASIX) tablet 20 mg 20 mg, oral, once, Starting S, For 1 Doses Administer post blood transfusion.	Every visit	S	Until discontin't'd