

CAGT- CLINIC PROCEDURE W CONSCIOUS SEDATION

version 10/10/2016

Provider Communication

	Interval	Defer Until	Duration
<input type="checkbox"/> Please place procedure, imaging, and referral orders using your preference list. These orders must be placed outside of the therapy plan to be scheduled. Details	Every 1 day	S	Until discont'd

Nursing Pre-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> Nursing communication Notify *** that the patient has arrived for procedure	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Notify the Marrow Technician that the patient has arrived	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Ascertain that informed consent has been obtained. If not, provider patient with preprinted consent form for marrow biopsy and aspiration for review. Do not begin procedure until consent has been obtained.	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Indication: ***. Risks: bleeding, infection, discomfort, damage to hip area.	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Identify adult who will accompany patient home after the procedure	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Have patient change into hospital gown	Once	S	1 treatment

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 25 mL/hr, continuous, Starting S Keep vein open	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9 % bolus 1,000 mL 1,000 mL, intravenous, at 500 mL/hr, for 120 Minutes, once, Starting S, For 1 Doses	Once	S	1 treatment
<input type="checkbox"/> morPHINE injection 4 mg, intravenous, once, Starting S, For 1 Doses	Once	S	1 treatment
<input type="checkbox"/> LORAZepam (ATIVAN) injection 1 mg, intravenous, every 4 hours PRN, sedation, Starting S	Once	S	1 treatment
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, every 6 hours PRN, itching, Starting S	Once	S	1 treatment

Intra-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> MIDAZolam (VERSED) injection 1-10 mg 1-10 mg, intravenous, PRN, sedation, Starting S	Once	S	1 treatment
<input type="checkbox"/> flumazenil (ROMAZICON) injection 0.3 mg 0.3 mg, intravenous, PRN, Sedation, Starting S	Once	S	1 treatment

Nursing Intra-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> Nursing communication During procedure: Monitor pulse and oxygen saturation continuously	Once	S	1 treatment
<input type="checkbox"/> Nursing communication During procedure: Record BP, pulse, oxygen saturation and respiratory rate every 5 minutes	Once	S	1 treatment
<input type="checkbox"/> Nursing communication During procedure: Monitor EKG continuously	Once	S	1 treatment

Nursing Post-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> Nursing communication After procedure: Record BP, pulse, oxygen saturation and respiratory rate every 15 minutes until discharge	Once	S	1 treatment
<input type="checkbox"/> Nursing communication After procedure: Perform and document Aldrete score every 30 minutes until discharge	Once	S	1 treatment
<input type="checkbox"/> Nursing communication After procedure: Patient to remain on unit for monitoring for at least 1 hour after procedure	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Provide post-procedure education when Aldrete score is 10	Once	S	1 treatment
<input type="checkbox"/> Nursing communication Discharge patient per guidelines and must be accompanied by an adult upon discharge	Once	S	1 treatment