Labs				
		Interval	Defer Until	Duration
	CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
	Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
	Magnesium level STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
	LDH STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
	Type and screen STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
Nursing C	orders			
		Interval	Defer Until	Duration
	Insert Peripheral IV or Access Portacath  Details	Every visit	S	Until discont'd
	Vital signs - T/P/R/BP Per Unit Protocol Details	Every visit	S	Until discont'd
	sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Every visit	S	Until discont'd
	HEParin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses ONLY IF PATIENT HAS PORTACATH: Flush portacath before removing ne	Every visit edle from portacath	S n.	Until discont'd
	sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
	Discontinue IV Details	Every visit	S	Until discont'd
	Discharge patient after completion of therapy, if stable Details	Every visit	S	Until discont'd
	Verify patient/caregiver received prescriptions and/or instruction as needed Details	Every visit	S	Until discont'd
Pre-Medic	eations			
		Interval	Defer Until	Duration
	ondansetron (ZOFRAN) injection 8 mg 8 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, PRN, mild pain (score 1-3), Starting S	Every visit	S	Until discont'd
IV Fluids				
		Interval	Defer Until	Duration
	sodium chloride 0.9 % 500 mL with magnesium sulfate infusion intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
	sodium chloride 0.9 % 1,000 mL with magnesium sulfate infusion intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
Supportiv	e Care			
		Interval	Defer Until	Duration
	ondansetron (ZOFRAN) injection 8 mg 8 mg, intravenous, every 6 hours PRN, nausea, vomiting, Starting S	Every visit	S	Until discont'd

	Interval	Defer Until	Duration
diphenhydrAMINE (BENADRYL) injection 25 mg	PRN	S	Until discont'd
25 mg, intravenous, once, Starting S, For 1 Doses			