

Labs

		Interval	Defer Until	Duration
<input type="checkbox"/> Epstein Barr Virus (EBV) by PCR	STAT, Status: Future, Expires: S+366, Clinic Collect Specimen Source: External send out- Draw and hold EBV DNA by PCR 1 purple top to Dr. Heslop lab at TCH	Every visit	S	Until discont'd

Nursing Orders

		Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details		Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Maintain IV access Details		Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses		Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S		Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details		Every visit	S	Until discont'd

Pre-Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.		Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses		Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 50-100 mg 50-100 mg, intravenous, Starting S Administer 30 minutes before infusion.		Every visit	S	Until discont'd

Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> immune globulin (GAMUNEX-C) 10 % solution 0.5 g/kg (Adjusted) 0.5 g/kg, intravenous, once, Starting H+30 Minutes, For 1 Doses	Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness.	Every 1 day	S	4 treatments

Hypersensitivity

		Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources).		PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S		PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, Starting S, For 1 Doses		PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, Starting S, For 1 Doses		PRN	S	Until discont'd
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mg, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses		PRN	S	Until discont'd

		Interval	Defer Until	Duration
<input type="checkbox"/>	hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/>	promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/>	sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discont'd