

PICC Line or Implanted Port

	Interval	Defer Until	Duration
<input type="checkbox"/> Change dressing: PICC Details	1 time a week	S	Until discont'd
<input type="checkbox"/> Change dressing: Port-a-cath Port-a-cath dressing is changed ONLY when non-coring needle is changed and as needed.	1 time a week	S	Until discont'd
<input type="checkbox"/> Access Portacath Routine, Clinic Performed	Every visit	S	Until discont'd
<input type="checkbox"/> Replace StatLock Details	1 time a week	S	Until discont'd
<input type="checkbox"/> Notify Physician if arm swelling, drainage, or other complications arise concerning the central line Details	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

Centrally Inserted Catheter or Temporary Dialysis Catheter

	Interval	Defer Until	Duration
<input type="checkbox"/> Change dressing Details	1 time a week	S	Until discont'd
<input type="checkbox"/> Notify Physician if arm swelling, drainage, or other complications arise concerning the central line Details	Every visit	S	Until discont'd

Line Flush

	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, every 8 hours, Starting S Aspirate and flush for saline locked IV. ONLY USE 10 mL SYRINGE OR LARGER TO FLUSH LINE.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 20 mL 20 mL, intra-catheter, every 8 hours, Starting S Flush Port-a-cath with 20 mL. ONLY USE 10 CC OR LARGER TO FLUSH LINE.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S Aspirate and flush 10 mL IV push before and after medications and as needed using turbulent method. ONLY USE 10 mL SYRINGE OR LARGER TO FLUSH LINE.	PRN	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 20 mL 20 mL, intravenous, PRN, line care, Starting S Flush with 20 mLs IV push after blood withdrawals and TPN. ONLY USE 10 mL SYRINGE OR LARGER TO FLUSH LINE.	PRN	S	Until discont'd

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> HEParin, porcine (PF) injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> alteplase (CATHFLO) for NON Quinton and NON Hemodialysis Catheters 1 mg 1 mg, intra-catheter, PRN, Starting S 1 mg dose of Cathflo is for NON Quinton and Hemodialysis Catheter Clearance. For occluded catheter port. Up to 2 doses may be used per port separated by 120 minutes.	PRN	S	Until discont'd
<input type="checkbox"/> alteplase (CATHFLO) for Quinton and Hemodialysis Catheters 2 mg	PRN	S	Until discont'd

Medications (continued)

	Interval	Defer Until	Duration
2 mg, intra-catheter, PRN, Starting S 2 mg dose of Cathflo is for Quinton and Hemodialysis Catheter Clearance. For occluded catheter port. Up to 2 doses may be used per port separated by 120 minutes.			

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC hemogram Routine, Status: Future, Expires: S+366, Clinic Collect	1 time a week	S	Until discontin'd
<input type="checkbox"/> Basic metabolic panel Routine, Status: Future, Expires: S+366, Clinic Collect	1 time a week	S	Until discontin'd
<input type="checkbox"/> Sedimentation rate Routine, Status: Future, Expires: S+366, Clinic Collect	1 time a week	S	Until discontin'd
<input type="checkbox"/> C-reactive protein Routine, Status: Future, Expires: S+366, Clinic Collect	1 time a week	S	Until discontin'd
<input type="checkbox"/> Blood culture, aerobic & anaerobic Routine, Status: Future, Expires: S+366, Clinic Collect	1 time a week	S	Until discontin'd
<input type="checkbox"/> Blood culture, fungus Routine, Status: Future, Expires: S+366, Clinic Collect, Blood	1 time a week	S	Until discontin'd