

## Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> Basic metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> Cytomegalovirus by PCR STAT, Status: Future, Expires: S+366, Clinic Collect Specimen Source:	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> Cytomegalovirus antigen STAT, Status: Future, Expires: S+366, Clinic Collect	Every 14 days	S	Until discontin't'd

## Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert peripheral IV Routine, Clinic Performed	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> Nursing communication Please verify patient received 2 gram dose of oral probenecid approximately 3 hours prior to initiation of cidofovir infusion.	Every 14 days	S	Until discontin't'd
<input type="checkbox"/> Nursing communication Please remind patient to take 1 gram (2 tablets) both 2 hours and 8 hours after end of infusion.	Every 14 days	S	Until discontin't'd

## Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> cidofovir (VISTIDE) 5 mg/kg in sodium chloride 0.9% 1,000 mL IVPB 5 mg/kg, intravenous, for 3 Hours, once, Starting H, For 1 Doses [cidofovir]Reason for Therapy:	Every 14 days	S	Until discontin't'd

## Discharge Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Discontinue IV Routine	Every 14 days	S	Until discontin't'd