

## Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Obtain vital signs every 30 minutes for first hour during hydration, then every 15 minutes during Cytoxan administration, then every 30 minutes for remainder of infusion.	Once	S	1 treatment
<input type="checkbox"/> Advise patient to have a light meal prior to chemotherapy, force fluid for 24-48 hours post chemotherapy, and empty bladder every 2 hours. Details	Once	S	1 treatment
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Once	S	1 treatment
<input type="checkbox"/> Discontinue IV Details	Once	S	1 treatment

## Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> ondansetron (ZOFTRAN) tablet 8 mg 8 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Once	S	1 treatment
<input type="checkbox"/> ondansetron (ZOFTRAN) IV 8 mg 8 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Once	S	1 treatment
<input type="checkbox"/> dexamethasone (DECADRON) IV 8 mg 8 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Once	S	1 treatment
<input type="checkbox"/> palonosetron (ALOXI) injection 0.25 mg 0.25 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Once	S	1 treatment
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Once	S	1 treatment

## Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> cyclophosphamide (CYTOXAN) 750 mg in sodium chloride 0.9 % 250 mL chemo IVPB 750 mg, intravenous, for 60 Minutes, once, Starting H+30 Minutes, For 1 Doses	Once	S	1 treatment

## Supportive Care

	Interval	Defer Until	Duration
<input type="checkbox"/> promethazine (PHENERGAN) injection 25-50 mg 25-50 mg, injection, every 4 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 4 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFTRAN) tablet 8 mg 8 mg, oral, every 8 hours PRN, nausea, vomiting, Nausea prevention, Starting S, For 2 Doses Zofran 8 mg Q8H x 2 doses, then 8 mg Q12H x 24-48 hours.	PRN	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFTRAN) tablet 8 mg 8 mg, oral, every 12 hours PRN, nausea, vomiting, Nausea prevention, Starting S, For 48 Hours Zofran 8 mg Q8H x 2 doses, then 8 mg Q12H x 24-48 hours.	PRN	S	Until discont'd

	Interval	Defer Until	Duration
<input type="checkbox"/> metoclopramide (REGLAN) tablet 10 mg 10 mg, oral, 3 times daily, Starting S, For 4 Days	PRN	S	Until discontin't'd
<input type="checkbox"/> dexamethasone (DECADRON) tablet 8 mg 8 mg, oral, every morning, Starting S+1, For 3 Doses Administer on Day 2-4.	PRN	S	Until discontin't'd
<input type="checkbox"/> mesna (MESNEX) tablet 800 mg 800 mg, oral, every 4 hours, Starting S, For 2 Doses Administer at 4 hours and 8 hours AFTER Mesna dose.	PRN	S	Until discontin't'd