

## Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential Routine, Status: Future, Expires: S+366, Clinic Collect, Prior to initiating Cytoxan, on Day 10 and as directed by the physician.	Every visit	S	Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel Routine, Status: Future, Expires: S+366, Clinic Collect, Prior to initiating Cytoxan, on Day 10 and as directed by the physician.	Every visit	S	Until discont'd
<input type="checkbox"/> Urinalysis, automated with microscopy Routine, Status: Future, Expires: S+366, Clinic Collect, Every visit.	Every visit	S	Until discont'd

## Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Obtain vital signs every 30 minutes for first hour during hydration, then every 15 minutes during Cytoxan administration, then every 30 minutes for remainder of infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> Advise patient to have a light meal prior to chemotherapy, force fluid for 24-48 hours post chemotherapy, and empty bladder every 2 hours. Details	Every visit	S	Until discont'd
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

## Hydration

	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride 0.45 % infusion 500 mL 500 mL, intravenous, at 500 mL/hr, for 30 Minutes, once, Starting S, For 1 Doses Infuse before and after Cytoxan at max rate of 500 mL/hr for hydration.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9 % infusion 500 mL 500 mL, intravenous, at 500 mL/hr, for 30 Minutes, once, Starting S, For 1 Doses Infuse before and after Cytoxan at max rate of 500 mL/hr for hydration.	Every visit	S	Until discont'd

## Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> ondansetron (ZOFTRAN) IV 8 mg 8 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> dexamethasone (DECADRON) IV 8 mg 8 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> palonosetron (ALOXI) injection 0.25 mg 0.25 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection intravenous, once, Starting S, For 1 Doses Administer 30 minutes prior to start of Cytoxan infusion.	Every visit	S	Until discont'd

## Medications

	Interval	Defer Until	Duration
--	----------	-------------	----------

**Medications (continued)**

	Interval	Defer Until	Duration
<input type="checkbox"/> cyclophosphamide (CYTOXAN) in sodium chloride 0.9 % 250 mL chemo IVPB intravenous, for 60 Minutes, once, Starting H+30 Minutes, For 1 Doses Administer every month x 6 months.	Day 1 of every 1 month	S	6 treatments
<input type="checkbox"/> mesna (MESNEX) 1,000 mg in sodium chloride 0.9 % 100 mL chemo IVPB 1,000 mg, intravenous, for 15 Minutes, once, Starting H+90 Minutes, For 1 Doses	Day 1 of every 1 month	S	6 treatments
<input type="checkbox"/> cyclophosphamide (CYTOXAN) in sodium chloride 0.9 % 250 mL chemo IVPB intravenous, for 60 Minutes, once, Starting H+30 Minutes, For 1 Doses Administer every 3 months x 18 months.	Day 1 of every 3 months	S+179	18 treatments
<input type="checkbox"/> mesna (MESNEX) 1,000 mg in sodium chloride 0.9 % 100 mL chemo IVPB 1,000 mg, intravenous, for 15 Minutes, once, Starting H+90 Minutes, For 1 Doses	Day 1 of every 3 months	S+179	18 treatments

**Supportive Care**

	Interval	Defer Until	Duration
<input type="checkbox"/> promethazine (PHENERGAN) injection 25-50 mg 25-50 mg, injection, every 4 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 4 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFTRAN) tablet 8 mg 8 mg, oral, every 8 hours PRN, nausea, vomiting, Nausea prevention, Starting S, For 2 Doses Zofran 8 mg Q8H x 2 doses, then 8 mg Q12H x 24-48 hours.	PRN	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFTRAN) tablet 8 mg 8 mg, oral, every 12 hours PRN, nausea, vomiting, Nausea prevention, Starting S, For 48 Hours Zofran 8 mg Q8H x 2 doses, then 8 mg Q12H x 24-48 hours.	PRN	S	Until discont'd
<input type="checkbox"/> metoclopramide (REGLAN) tablet 10 mg 10 mg, oral, 3 times daily, Starting S, For 4 Days	PRN	S	Until discont'd
<input type="checkbox"/> dexamethasone (DECADRON) tablet 8 mg 8 mg, oral, every morning, Starting S+1, For 3 Doses Administer on Day 2-4.	PRN	S	Until discont'd
<input type="checkbox"/> mesna (MESNEX) tablet 800 mg 800 mg, oral, every 4 hours, Starting S, For 2 Doses Administer at 4 hours and 8 hours after Mesna IV dose.	PRN	S	Until discont'd