CYTOMEGALOVIRUS IMMUNE GLOBULIN (CYTOGAM) INFUSION

Labs					
		Basic metabolic panel	Interval Every visit	Defer Until S	Duration Until discont'd
	_	STAT, Status: Future, Expires: S+366, Clinic Collect	,		
		CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
		Hepatic function panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
Nursin	g O	rders			
			Interval	Defer Until	Duration
		Vital signs - T/P/R/BP Per Unit Protocol Vital Signs PRE & POST Infusion.	Every visit	S	Until discont'd
		Nursing communication: Maintain IV access Details	Every visit	S	Until discont'd
		sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd
		sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
		Discontinue IV Details	Every visit	S	Until discont'd
Pre-Me	edic	ations			
			Interval	Defer Until	Duration
		diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
		acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
		methylPREDNISolone sodium succinate (Solu-MEDROL) 250 mg in sodium chloride 0.9% 100 mL IVPB 250 mg, intravenous, for 30 Minutes, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
Loading Dose					
			Interval	Defer Until	Duration
		cytomegalovirus immune globulin (CYTOGAM) IVPB 250 mg/kg (Dosing Weight) 250 mg/kg, intravenous, once, Starting S, For 1 Doses Per manufacturer's guidelines, according to patient tolerance.	Every 2 weeks	S	4 treatments
Mainte	nan	nce Dose			
			Interval	Defer Until	Duration
		cytomegalovirus immune globulin (CYTOGAM) IVPB 250 mg/kg (Dosing Weight) 250 mg/kg, intravenous, once, Starting S, For 1 Doses Per manufacturer's guidelines, according to patient tolerance.	Every 4 weeks	S+76	Until discont'd