rders			
	Interval	Defer Until	Duration
Vital signs - T/P/R/BP Per Unit Protocol	Every visit	S	Until discont'd
Instruct patient to report symptoms of chills, fever or pain.			
Nursing communication: Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered Details	Every visit	S	Until discont'o
Treatment conditions	Every visit	S	Until discont'd
Notify provider if Ca+ level is LESS than or EQUAL to 8.5			
	Interval	Defer Until	Duration
Ionized calcium	Every visit	S	Until discont'd
Routine, Status: Future, Expires: S+366, Clinic Collect	-		
Comprehensive metabolic panel	Every visit	S	Until discont'd
Routine, Status: Future, Expires: S+366, Clinic Collect	,		
POC pregnancy, urine	Every visit	S	Until discont'd
Routine, Point Of Care	•		
ns			
	Interval	Defer Until	Duration
denosumab (PROLIA) syringe 60 mg	Day 1 of every 6	S	2 treatments
60 mg, subcutaneous, once, Starting S, For 1 Doses			
sitivity			
Sitivity			
Sitivity	Interval	Defer Until	Duration
acetaminophen (TYLENOL) tablet 650 mg	Interval PRN	Defer Until S	Duration Until discont'd
		_	
acetaminophen (TYLENOL) tablet 650 mg		_	
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	PRN	S	Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S	PRN	S	Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg	PRN PRN PRN	S	Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S,	PRN PRN PRN For 1 Doses	S S	Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg	PRN PRN PRN For 1 Doses PRN	S	Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S	PRN PRN PRN For 1 Doses PRN S, For 1 Doses	S S S	Until discont'd Until discont'd Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg	PRN PRN PRN For 1 Doses PRN	S S	Until discont'd Until discont'd Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN PRN PRN For 1 Doses PRN S, For 1 Doses PRN	S S S	Until discont'd Until discont'd Until discont'd Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg	PRN PRN PRN For 1 Doses PRN S, For 1 Doses PRN PRN	S S S	Until discont'd Until discont'd Until discont'd Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN PRN PRN For 1 Doses PRN S, For 1 Doses PRN PRN	S S S S	Until discont'd Until discont'd Until discont'd Until discont'd Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses promethazine (PHENERGAN) tablet 25 mg	PRN PRN PRN For 1 Doses PRN S, For 1 Doses PRN PRN	S S S	Until discont'd Until discont'd Until discont'd Until discont'd Until discont'd Until discont'd
acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN PRN PRN For 1 Doses PRN S, For 1 Doses PRN PRN	S S S S	Until discont'd Until discont'd
	Vital signs - T/P/R/BP Per Unit Protocol Instruct patient to report symptoms of chills, fever or pain. Nursing communication: Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered Details Treatment conditions Notify provider if Ca+ level is LESS than or EQUAL to 8.5 Ionized calcium Routine, Status: Future, Expires: S+366, Clinic Collect Comprehensive metabolic panel Routine, Status: Future, Expires: S+366, Clinic Collect POC pregnancy, urine Routine, Point Of Care denosumab (PROLIA) syringe 60 mg	Vital signs - T/P/R/BP Per Unit Protocol Instruct patient to report symptoms of chills, fever or pain. Nursing communication: Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered Details Treatment conditions Notify provider if Ca+ level is LESS than or EQUAL to 8.5 Interval Every visit Interval Every visit Routine, Status: Future, Expires: S+366, Clinic Collect Comprehensive metabolic panel Routine, Status: Future, Expires: S+366, Clinic Collect POC pregnancy, urine Routine, Point Of Care ns Interval Every visit Every visit Interval Every visit Every visit Every visit For pregnancy, urine Routine, Point Of Care ns Interval Day 1 of every 6 months for months	Vital signs - T/P/R/BP Per Unit Protocol Instruct patient to report symptoms of chills, fever or pain. Nursing communication: Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered Details Treatment conditions Notify provider if Ca+ level is LESS than or EQUAL to 8.5 Interval Defer Until Every visit S