.abs				
		Interval	Defer Until	Duration
	CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
	Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
	hCG qualitative, urine screen STAT, Status: Future, Expires: S+366, Clinic Collect Is this the first void of the morning?	Every visit	S	Until discont'd
	Ionized calcium STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
lursing (Orders			
		Interval	Defer Until	Duration
	Vital signs - T/P/R/BP Per Unit Protocol Instruct patient to report symptoms of chills, fever or pain.	Every visit	S	Until discont'd
	Nursing communication: Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered Details	Every visit	S	Until discont'd
	Treatment conditions Notify provider if Ca+ level is LESS than or EQUAL to 8.5	Every visit	S	Until discont'd
/ledicatio	ons			
	denosumab (XGEVA) injection 120 mg	Interval Every 4 weeks	Defer Until S	Duration Until discont'd
	120 mg, subcutaneous, once, Starting S, For 1 Doses			
lyperser	sitivity			
71		Interval	Defer Until	Duration
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	PRN	S	Until discont'd
	diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S	PRN	S	Until discont'd
	diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S	PRN , For 1 Doses	S	Until discont'd
	diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting 9	PRN S, For 1 Doses	S	Until discont'd
	or mg, mark once, once it is in the might be determined in the control of the might			
	epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'd
	epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg	PRN	S	Until discont'd Until discont'd
	epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg	PRN		