

FERRIC GLUCONATE (FERRLECIT) INFUSION

version 10/4/2018

Labs

| | Interval | Defer Until | Duration |
|---|-------------|-------------|-----------------|
| <input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect | Every visit | S | Until discont'd |

Nursing Orders

| | Interval | Defer Until | Duration |
|--|-------------|-------------|-----------------|
| <input type="checkbox"/> Insert Peripheral IV or Access Portacath Details | Every visit | S | Until discont'd |
| <input type="checkbox"/> For new patients, patients requiring pre-medication, and patients with previous reactions to IV iron, monitor 1 hour following infusion for shortness of breath, wheezing, hypotension, tachycardia, or hives. Details | Every visit | S | Until discont'd |
| <input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S | Every visit | S | Until discont'd |
| <input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S | Every visit | S | Until discont'd |
| <input type="checkbox"/> Discontinue IV Details | Every visit | S | Until discont'd |

Pre-Medications

| | Interval | Defer Until | Duration |
|--|-------------|-------------|-----------------|
| <input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion. | Every visit | S | Until discont'd |
| <input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25-50 mg 25-50 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion. | Every visit | S | Until discont'd |
| <input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 50-100 mg 50-100 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion. | Every visit | S | Until discont'd |
| <input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion. Not covered for medicare patients. | Every visit | S | Until discont'd |
| <input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg 180 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion. | Every visit | S | Until discont'd |

Medications

| | Interval | Defer Until | Duration |
|--|----------|-------------|-------------|
| <input type="checkbox"/> ferric gluconate (FERRLECIT) 125 mg in sodium chloride 0.9% 100 mL IVPB 125 mg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses Not to exceed 2.1 mg/min (duration = 60 min) | Once | S | 1 treatment |

Supportive Care

| | Interval | Defer Until | Duration |
|--|-------------|-------------|-----------------|
| <input type="checkbox"/> promethazine (PHENERGAN) injection 25 mg 25 mg, injection, every 4 hours PRN, nausea, vomiting, Starting S | Every visit | S | Until discont'd |

Emergency Medications

| | Interval | Defer Until | Duration |
|---|----------|-------------|-----------------|
| <input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg | PRN | S | Until discont'd |

| | Interval | Defer Until | Duration |
|--|----------|-------------|-----------------|
| 100 mg, intravenous, once, Starting S, For 1 Doses | | | |
| <input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses | PRN | S | Until discont'd |
| <input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once, Starting S, For 1 Doses | PRN | S | Until discont'd |