

**Labs**

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Sedimentation rate STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> C-reactive protein STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd

**Nursing Orders**

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Details	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Review lab results for Hep B, Hep C, and TB Details	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for signs of infusion reaction. For MINOR INFUSION REACTIONS: fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea. Take the following steps: Stop or SLOW infusion rate by 50%. If symptoms do not resolve within 5-10 minutes, give supportive/emergency medications as ordered, and notify MD. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor closely until completion.	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of infusion reaction. For SEVERE INFUSION REACTIONS: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension. Take the following steps: STOP INFUSION, and start Normal Saline IV bolus (as ordered), Notify physician and administer emergency medications (as ordered). Monitor vital signs. Consult physician for further orders.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

**Pre-Medications**

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 500 mg 500 mg, oral, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

**Medications**

	Interval	Defer Until	Duration
<input type="checkbox"/> golimumab (SIMPONI ARIA) 2 mg/kg in sodium chloride 0.9% 100 mL IVPB 2 mg/kg, intravenous, for 30 Minutes, once, Starting H+30 Minutes, For 1 Doses Use an infusion set with an in-line, sterile, non-pyrogenic filter (0.22 micrometer or less). Do not infuse other agents in the same IV line. Administer on Week 0 and 4.	Every 4 weeks	S	2 treatments
<input type="checkbox"/> golimumab (SIMPONI ARIA) 2 mg/kg in sodium chloride 0.9% 100 mL IVPB 2 mg/kg, intravenous, for 30 Minutes, once, Starting H+30 Minutes, For 1 Doses Use an infusion set with an in-line, sterile, non-pyrogenic filter (0.22 micrometer or less). Do not infuse other agents in the same IV line. Administer on Week 12 and every 8 weeks thereafter.	Every 8 weeks	S+83	Until discont'd

## Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discont'd