Labs	;						
			Interval	Defer Until	Duration		
		Hepatitis B surface antigen STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd		
		Hepatitis B surface antibody STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd		
Nursing Orders							
			Interval	Defer Until	Duration		
		Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd		
		Nursing communication: Contact physician and discontinue HBIG infusion if the patient experiences any of the following symptoms: hypotension, tachycardia, severe myalgias, shortness of breath Details	Every visit	S	Until discont'd		
		sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd		
		HEParin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses ONLY IF PATIENT HAS PORTACATH: Flush portacath before removing need	Every visit	S	Until discont'd		
		In Rare Event of Anaphylaxis: **Stop HBIG infusion immediately **Administer 0.3-0.5 mg = 0.3-0.5 mL of the 1:1000 epinephrine subcutaneously **Notify attending physician for advisement immediately Details	Every visit	S	Until discont'd		
		sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd		
		Discontinue IV Details	Every visit	S	Until discont'd		
Pre-Medications							
			Interval	Defer Until	Duration		
		acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd		
		diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd		
Medi	catio	ns	_				
		hepatitis B immune globulin (HEPAGAM B) 9,360 Units in sodium chloride 0.9% 250 mL IVPB 9,360 Units, intravenous, for 3 Hours, once, Starting H+30 Minutes, For 1 Dos	Interval Every 4 weeks es	Defer Until S	Duration Until discont'd		
Eme	rgenc	y Medications					
			Interval	Defer Until	Duration		
		epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.3-0.5 mg 0.3-0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'd		
		diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'd		
		hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'd		