

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Sedimentation rate STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> C-reactive protein STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Zinc level, serum STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Ferritin level STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Iron level STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Total iron binding capacity STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Vitamin D 25 hydroxy level STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Vitamin B12 level STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Vitamin A level, plasma or serum STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Folate level STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> TB GOLD Quantiferon STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Details	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Observe patient for 30 minutes Details	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for any signs of infusion reaction. For MINOR INFUSION REACTIONS (fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea) stop or slow infusion rate by 50%. If symptoms do not resolve within 5-10 minutes give Benadryl as ordered. Notify MD if symptoms do not resolve within 5-10 minutes. With MD approval, give Solu-medrol as ordered. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor patient closely until completion.	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of SEVERE INFUSION REACTION: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension. For severe infusion reaction take the following steps; STOP Infusion and start Normal Saline solution, Notify physician and administer Benadryl as ordered, Monitor vital signs, Administer Epinephrine as ordered. Consult MD for further orders.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> HEParin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses ONLY IF PATIENT HAS PORTACATH: Flush portacath before removing needle from portacath.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL	Every visit	S	Until discont'd

Nursing Orders (continued)

	Interval	Defer Until	Duration
250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S			
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 500 mg 500 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> famotidine (PEPCID) tablet 40 mg 40 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg 180 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 125 mg 125 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	PRN	S	Until discont'd

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> inFLIXimab (REMICADE) 5 mg/kg in sodium chloride 0.9% 250 mL IVPB 5 mg/kg, intravenous, for 120 Minutes, once, Starting S, For 1 Doses Use an in-line, sterile, non-pyrogenic, low protein-binding filter with 1.2 micron pore size or less. Infuse at 10 mL/hour x 15 minutes, then infuse at 20 mL/hour x 15 minutes, then infuse at 40 mL/hour x 15 minutes, then infuse at 80 mL/hour x 15 minutes, then infuse at 150 mL/hour x 30 minutes, then infuse at 250 mL/hour until complete.	Every 8 weeks	S	Until discont'd

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours PRN, mild pain (score 1-3), headaches, fever, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for mild to moderate infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd

	Interval	Defer Until	Duration
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> ondansetron (ZOFTRAN) injection 4 mg 4 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discontinuation