

**Labs**

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Ferritin level STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Total iron binding capacity STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd

**Nursing Orders**

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Observe 1 hour and check vital signs after each dose.	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for signs of *MINOR INFUSION REACTION*: fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea. For minor infusion reaction take the following steps: stop or slow infusion rate by 50%. If symptoms do not resolve within 5-10 minutes give Benadryl as ordered. Notify MD if symptoms do not resolve within 5-10 minutes. With MD approval, give Solu-medrol as ordered. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor patient closely until completion.	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of *SEVERE INFUSION REACTION*: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension. For severe infusion reaction take the following steps; STOP Infusion and start normal saline solution. Notify physician and administer Benadryl as ordered. Monitor vital signs. Administer Epinephrine as ordered. Consult MD for further orders.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

**Hydration**

	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 50 mL/hr, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

**Medications**

	Interval	Defer Until	Duration
<input type="checkbox"/> iron dextran (INFED) 25 mg in sodium chloride 0.9% 25 mL IVPB 25 mg, intravenous, for 15 Minutes, once, Starting S, For 1 Doses Test dose; Observe 1 hour and check vital signs.	Once	S	1 treatment
<input type="checkbox"/> iron dextran (INFED) in sodium chloride 0.9% 100 mL IVPB intravenous, for 30 Minutes, once, Starting H+1 Hours, For 1 Doses Do not exceed 50 mg/minute; Observe 1 hour and check vital signs.	Once	S	1 treatment

**Hypersensitivity**

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours PRN, mild pain (score 1-3), headaches, fever, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for mild to moderate infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mg, subcutaneous, once PRN, anaphylaxis, infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd