

Nursing Orders

		Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details		Every visit	S	Until discont'd
<input type="checkbox"/> For new patients, patients requiring pre-medication, and patients with previous reactions to IV iron, monitor 1 hour following infusion for shortness of breath, wheezing, hypotension, tachycardia, or hives. Details		Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S		Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S		Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details		Every visit	S	Until discont'd

Pre-Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.		Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 25-50 mg 25-50 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.		Every visit	S	Until discont'd
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 50-100 mg 50-100 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.		Every visit	S	Until discont'd
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion. Not covered for medicare patients.		Every visit	S	Until discont'd
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg 180 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.		Every visit	S	Until discont'd

Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> iron sucrose (VENOFER) 200 mg in sodium chloride 0.9% 100 mL IVPB 200 mg, intravenous, for 30 Minutes, once, Starting S, For 1 Doses		Every 3 weeks	S	5 treatments

Supportive Care

		Interval	Defer Until	Duration
<input type="checkbox"/> promethazine (PHENERGAN) injection 25 mg 25 mg, intravenous, every 4 hours PRN, nausea, vomiting, Starting S		Every visit	S	Until discont'd

Emergency Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, infusion reaction, Starting S, For 1 Doses		PRN	S	Until discont'd
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, Starting S, For 1 Doses		PRN	S	Until discont'd
<input type="checkbox"/> epinephrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses		PRN	S	Until discont'd

