

IVIG BOOSTER INFUSION (ST. JOHNS - DEFAULT PT SUPPLIED)

version 1/30/2018

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	1 time a week	S	Until discontinuation

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	1 time a week	S	Until discontinuation
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Vital Signs PRE & POST Infusion.	1 time a week	S	Until discontinuation
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	1 time a week	S	Until discontinuation
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	1 time a week	S	Until discontinuation
<input type="checkbox"/> Discontinue IV Details	1 time a week	S	Until discontinuation

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 12.5 mg 12.5 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	1 time a week	S	Until discontinuation
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	1 time a week	S	Until discontinuation

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> IMMUNE GLOBULIN (IGG) 10% ORDERABLE FOR THERAPY PLANS 0.5 g/kg, intravenous, once, Starting H+30 Minutes, For 1 Doses Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness. DO NOT SUBSTITUTE:	1 time a week	S	Until discontinuation

Post-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> Heparin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, PRN, line care, Starting S	PRN	S	Until discontinuation