

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> Basic metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Hepatic function panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Creatinine level STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> BUN level STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd
<input type="checkbox"/> Urinalysis, automated with microscopy STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'd

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Vital Signs PRE & POST Infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Maintain IV access Details	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) 250 mg in sodium chloride 0.9% 100 mL IVPB 250 mg, intravenous, for 30 Minutes, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> IMMUNE GLOBULIN (IGG) 10% ORDERABLE FOR THERAPY PLANS 0.5 g/kg (Adjusted) 0.5 g/kg, intravenous, once, Starting S, For 1 Doses This IVIG entry is for Day 1. Nurses: please release this entry for Day 1.	Every 4 weeks	S	Until discont'd

Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness.
DO NOT SUBSTITUTE:

Medications (continued)

	Interval	Defer Until	Duration
<input type="checkbox"/> IMMUNE GLOBULIN (IGG) 10% ORDERABLE FOR THERAPY PLANS 0.5 g/kg (Adjusted) 0.5 g/kg, intravenous, once, Starting S, For 1 Doses This IVIG entry is for Day 2 Nurses: please release this entry for Day 2. Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness. DO NOT SUBSTITUTE:	Every 4 weeks	S	Until discontinued
<input type="checkbox"/> IMMUNE GLOBULIN (IGG) 10% ORDERABLE FOR THERAPY PLANS 0.5 g/kg (Adjusted) 0.5 g/kg, intravenous, once, Starting S, For 1 Doses This IVIG entry is for Day 3. Nurses: please release this entry for Day 3. Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness. DO NOT SUBSTITUTE:	Every 4 weeks	S	Until discontinued
<input type="checkbox"/> IMMUNE GLOBULIN (IGG) 10% ORDERABLE FOR THERAPY PLANS 0.5 g/kg (Adjusted) 0.5 g/kg, intravenous, once, Starting S, For 1 Doses This IVIG entry is for Day 4. Nurses: please release this entry for Day 4. Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness. DO NOT SUBSTITUTE:	Every 4 weeks	S	Until discontinued

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours PRN, mild pain (score 1-3), Starting S	PRN	S	Until discontinued
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for mild to moderate infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinued
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinued
<input type="checkbox"/> epinephrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mg, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinued
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinued
<input type="checkbox"/> methylprednisolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinued
<input type="checkbox"/> ondansetron (ZOFRAN) injection 4 mg 4 mg, intramuscular, once PRN, nausea, vomiting, Starting S, For 1 Doses	PRN	S	Until discontinued
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discontinued
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discontinued