

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication: Maintain IV access Details	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection intravenous, Starting S Administer 30 minutes before infusion.	Every visit	S	Until discont'd

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> immune globulin (GAMUNEX-C) 10 % solution 1 g/kg (Adjusted) 1 g/kg, intravenous, once, Starting H+30 Minutes, For 1 Doses Initiate infusion at 30 mL/hr for 60 minutes then double rate every 60 minutes as tolerated to a goal rate of 120 mL/hr. The patient is not tolerating the infusion if they develop headache, flushing, itching, rash, nausea & vomiting, muscle pain, tachycardia (heart rate >100 BPM), a temperature elevation of 1.8 degrees Fahrenheit, hypotension (SBP < 90 mmHg), hypertension (SBP > 180 mmHg), chills, problems breathing, or chest tightness.	Every 1 day	S	2 treatments