LEUPROLIDE (ELIGARD) SQ INJECTION

		Interval	Defer Until	Duration
	hCG qualitative, urine screen STAT, Status: Future, Expires: S+366, Clinic Collect Is this the first void of the morning? Prior to initiating therapy	Once	S	1 treatment
	Prostate specific antigen STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discont'c
icatio	ns			
		Interval	Defer Until	Duration
	leuprolide (ELIGARD) injection 7.5 mg	Every 4 weeks	S	Until discont'o
	7.5 mg, subcutaneous, once, Starting S, For 1 Doses			
	leuprolide (3 month) (ELIGARD) injection 22.5 mg	Every 12 weeks	S	Until discont'o
	22.5 mg, subcutaneous, once, Starting S, For 1 Doses			
	leuprolide (4 month) (ELIGARD) injection 30 mg	Every 16 weeks	S	Until discont'o
	30 mg, subcutaneous, once, Starting S, For 1 Doses			
	leuprolide (6 month) (ELIGARD) injection 45 mg	Every 24 weeks	S	Until discont'o
	45 mg subcutaneous once Starting S. For 1 Doses			

45 mg, subcutaneous, once, Starting S, For 1 Doses