

## Pre Intrathecal Injection Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> NPO after midnight Except sips with meds.	Once	S	1 treatment
<input type="checkbox"/> Routine vital signs, neuro checks, O2 sats, and I&O's Details	Once	S	1 treatment
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Once	S	1 treatment
<input type="checkbox"/> Nursing communication: Maintain IV access Details	Once	S	1 treatment
<input type="checkbox"/> Please have patient/family sign consent for Intrathecal Trial Procedure: Intrathecal morphine injection trial Diagnosis/Condition: Physician:	Once	S	1 treatment
<input type="checkbox"/> Lumbar puncture tray to bedside Routine, Normal	Once	S	1 treatment
<input type="checkbox"/> Nursing communication: please have the following items available for intrathecal injection: Iodine swabs, (2) 3 cc sterile syringes, 7 1/2 and 6 1/2 sterile gloves, 4x4 gauze, Bandaid	Once	S	1 treatment
<input type="checkbox"/> Consult to PT eval and treat Routine, Hospital Performed, Status: Future, Expires: S+366, Special Instructions: Please evaluate patient prior to baclofen injection and post injection.	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Once	S	1 treatment
<input type="checkbox"/> HEParin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses ONLY IF PATIENT HAS PORTACATH: Flush portacath before removing needle from portacath.	Once	S	1 treatment
<input type="checkbox"/> Discontinue IV Details	Once	S	1 treatment

## Intrathecal Injections

	Interval	Defer Until	Duration
<input type="checkbox"/> morPHINE PF 1 mg/mL injection 1 mg 1 mg, intrathecal, once, Starting S, For 1 Doses For intrathecal use ONLY.	Once	S	1 treatment

## Post Intrathecal Injection Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Nursing communication Continue all active admission orders including pulse ox. Flat in bed until ***. Regular diet after ***. Pt to take home medications. Discharge patient home after last PT visit.	Once	S	1 treatment
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	Once	S	1 treatment
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S	Once	S	1 treatment
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, every 6 hours PRN, itching, Starting S	Once	S	1 treatment
<input type="checkbox"/> ondansetron (ZOFran) injection 4 mg	Once	S	1 treatment

	Interval	Defer Until	Duration
4 mg, intravenous, every 4 hours PRN, nausea, vomiting, Starting S			
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	Once	S	1 treatment
<input type="checkbox"/> promethazine (PHENERGAN) injection 25 mg 25 mg, intravenous, every 6 hours PRN, nausea, vomiting, Starting S	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, once, Starting S, For 1 Doses TKO during recovery	Once	S	1 treatment