

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Nursing communication Repeat regimen daily for *** days	Every visit	S	Until discont'd
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication DHE Infusion: o Monitor patient for side effects. Decrease the infusion rate and/or temporarily stop the infusion. If symptoms improve, continue infusion as best tolerated rate and dose. o Need recent NEGATIVE pregnancy test o Need recent EKG on file	Every visit	S	Until discont'd

Steroids

	Interval	Defer Until	Duration
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) IVPB 1,000 mg 1,000 mg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> dexamethasone (DECADRON) injection 4 mg 4 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

Antiepileptic

	Interval	Defer Until	Duration
<input type="checkbox"/> valproate sodium (DEPACon) 1,000 mg 1,000 mg, intravenous, for 60 Minutes, once, Starting H, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> levETIRAcetam (KEPPRA) IV 1,000 mg 1,000 mg, intravenous, once, Starting H, For 1 Doses	Every visit	S	Until discont'd

Miscellaneous

	Interval	Defer Until	Duration
<input type="checkbox"/> hydrOXYzine (ATARAX) tablet 25 mg 25 mg, oral, once, Starting H, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> LORAZepam (ATIVAN) injection 1 mg 1 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> keTOROlac (TORadol) injection 30 mg 30 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> magnesium sulfate IVPB 1 g 1 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9 % bolus 1,000 mL 1,000 mL, intravenous, for 60 Minutes, once, Starting S, For 1 Doses For DAY 1 only	Once	S	1 treatment

Vasoconstrictive Agents

	Interval	Defer Until	Duration
<input type="checkbox"/> SUMATriptan succinate (IMITREX) injection 6 mg	Every visit	S	Until discont'd

Vasoconstrictive Agents (continued)

	Interval	Defer Until	Duration
6 mg, subcutaneous, once, Starting S, For 1 Doses Do not give within 24 hours of DHE or other ergotamine use. Do not use with other triptan medications. For subcutaneous administration only.			

Dopamine Antagonists

	Interval	Defer Until	Duration
<input type="checkbox"/> promethazine (PHENERGAN) IV syringe 12.5 mg 12.5 mg, intravenous, for 10 Minutes, once, Starting S, For 1 Doses	Every visit	S	Until discontin'td
<input type="checkbox"/> metoclopramide (REGLAN) injection 10 mg 10 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discontin'td
<input type="checkbox"/> chlorproMAZINE (THORAZINE) 12.5 mg in sodium chloride 0.9 % 50 mL IVPB 12.5 mg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses [chlorproMAZINE]Indication:	Every visit	S	Until discontin'td
<input type="checkbox"/> prochlorperazine (COMPAZINE) injection 10 mg 10 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discontin'td

DHE Infusion

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses Pre-med for DHE. Administer 30 minutes prior to start of DHE infusion.	Every visit	S	Until discontin'td
<input type="checkbox"/> ondansetron (ZOFTRAN) injection 4 mg 4 mg, intravenous, once, Starting S, For 1 Doses Pre-med for DHE. Administer 30 minutes prior to start of DHE infusion.	Every visit	S	Until discontin'td
<input type="checkbox"/> dihydroergotamine (DHE) 1 mg in sodium chloride 0.9 % 250 mL IVPB 1 mg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses If patient experiences side effects, decrease the infusion rate and/or temporarily stop the infusion. If symptoms improve, continue infusion as best tolerated rate and dose. Do not give within 24 hours of triptan use.	Every visit	S	Until discontin'td

PRN Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> ondansetron (ZOFTRAN) injection 4 mg 4 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discontin'td