

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Vital signs - T/P/R/BP Routine, Clinic Performed	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush Bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> HEParin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses ONLY IF PATIENT HAS PORTACATH: Flush portacath before removing needle from portacath	Every visit	S	Until discont'd
<input type="checkbox"/> Nursing communication Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, intravenous, once, Starting S, For 1 Doses Dilute with 10 mL of 0.9% sodium chloride injection and give IV push over at least 2 minutes	Every visit	S	Until discont'd
<input type="checkbox"/> dexamethasone (DECADRON) IV 10 mg 10 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> patisiran (lipid complex) (ONPATTRO) 30 mg in sodium chloride 0.9% 200 mL IVPB 30 mg, intravenous, once, Starting H+30 Minutes, For 1 Doses For patients GREATER than or EQUAL to 100 Kg. Initial infusion rate of 1 mL/minute for 15 minutes, then increase to ~3 mL/minute for the remainder of the infusion. Use a dedicated line with an infusion set containing a 1.2 micron PES in-line infusion filter. Use DEHP-free infusion sets and lines. Administer only through a free-flowing venous access line. Monitor the infusion site for infiltration. Manage extravasation according to local standard practice for non-vesicants. Onpattro™ (patisiran) is restricted to use in patients with prior financial approval in the outpatient setting. Do you attest that the patient is being treated in an approved care setting?	Every 3 weeks	S	Until discont'd
<input type="checkbox"/> patisiran (lipid complex) (ONPATTRO) 0.3 mg/kg in sodium chloride 0.9% 200 mL IVPB 0.3 mg/kg, intravenous, once, Starting H+30 Minutes, For 1 Doses For patients LESS than 100 Kg Initial infusion rate of 1 mL/minute for 15 minutes, then increase to ~3 mL/minute for the remainder of the infusion. Use a dedicated line with an infusion set containing a 1.2 micron PES in-line infusion filter. Use DEHP-free infusion sets and lines. Administer only through a free-flowing venous access line. Monitor the infusion site for infiltration. Manage extravasation according to local standard practice for non-vesicants.	Every 3 weeks	S	Until discont'd

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg	PRN	S	Until discont'd

	Interval	Defer Until	Duration
25 mg, oral, every 6 hours PRN, itching, Starting S			
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mg, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discontin'td
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, for hydration, Starting S, For 1 Doses	PRN	S	Until discontin'td