

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discontin'td
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discontin'td
<input type="checkbox"/> Uric acid level STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discontin'td

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discontin'td
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Obtain vital signs every 15 minutes throughout infusion and hydration. Monitor temperature every 30 minutes during infusion. Observe patient for 60 minutes following infusion. Instruct patient to report symptoms of chills, SOB, itching, or pain.	Every visit	S	Until discontin'td
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for signs of infusion reaction. For MINOR INFUSION REACTIONS: fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea. Take the following steps: Stop or SLOW infusion rate by 50%. If symptoms do not resolve within 5-10 minutes, give supportive/emergency medications as ordered, and notify MD. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor closely until completion.	Every visit	S	Until discontin'td
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of infusion reaction. For SEVERE INFUSION REACTIONS: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension, Take the following steps: STOP INFUSION, and start Normal Saline IV bolus (as ordered), Notify physician and administer emergency medications (as ordered). Monitor vital signs. Consult physician for further orders.	Every visit	S	Until discontin'td
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discontin'td
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discontin'td
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discontin'td

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 1,000 mg 1,000 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discontin'td
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) capsule 50 mg 50 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discontin'td
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 125 mg 125 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discontin'td

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> pegloticase (KRYSTEXXA) 8 mg in sodium chloride 0.9% 250 mL IVPB 8 mg, intravenous, for 120 Minutes, once, Starting S, For 1 Doses	Every 2 weeks	S	26 treatments

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S	PRN	S	Until discontin'td

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S	PRN	S	Until discontin'td
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontin'td
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discontin'td
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discontin'td