

Provider Communication

		Interval	Defer Until	Duration
<input type="checkbox"/> Contraindicated in patients with asthma.		Every 4 weeks	S	Until discont'd

Details

Pre-Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> albuterol (PROVENTIL) nebulizer solution 2.5 mg		Every 4 weeks	S	Until discont'd
	2.5 mg, nebulization, once, Starting S, For 1 Doses Administer via HHN. Aerosol Delivery Device: Hand-Held Nebulizer			
<input type="checkbox"/> levalbuterol (XOPENEX) nebulizer solution 0.63 mg		Every 4 weeks	S	Until discont'd
	0.63 mg, nebulization, once, Starting S, For 1 Doses Administer via HHN. Aerosol Delivery Device: Hand-Held Nebulizer			

Medications

		Interval	Defer Until	Duration
<input type="checkbox"/> pentamidine (PENTAM) inhalation solution 300 mg		Every 4 weeks	S	Until discont'd
	300 mg, inhalation, once, Starting S, For 1 Doses Administer via Respircard II nebulizer. May cause Q-T interval prolongation.			
<input type="checkbox"/> water for injection, sterile (PF) 6 mL		Every 4 weeks	S	Until discont'd
	6 mL, Other, once, Starting S, For 1 Doses Used to mix with Pentamidine powder.			