		Interval	Defer Until	Duration
	Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discor
	Vital signs - T/P/R/BP Per Unit Protocol Details	Every visit	S	Until discon
	Nursing communication: Observe patient for 30 minutes Details	Every visit	S	Until discon
	Nursing communication: Monitor patient for signs of infusion reaction. If patient has infusion reaction, stop infusion, notify physician, and administer emergency medications as ordered Details	Every visit	S	Until discon
	sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Every visit	S	Until discor
	sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discor
	Discontinue IV Details	Every visit	S	Until discor
catio	ns			
		Interval	Defer Until	Duration
	pentamidine (PENTAM) 4 mg/kg in dextrose 5% 250 mL IVPB	Every 1 day	S	Until discor
	4 mg/kg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses			
rsen	4 mg/kg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses sitivity			
rsen		 Interval	Defer Until	Duration
rsen		Interval PRN	Defer Until	Duration
	sitivity acetaminophen (TYLENOL) tablet 650 mg		_	Duration Until discor
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg	PRN PRN	S S	Duration Until discor Until discor
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg	PRN PRN PRN For 1 Doses PRN	S	Duration Until discor Until discor
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg	PRN PRN PRN For 1 Doses PRN	S S	Duration Until discor Until discor Until discor
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 1 mg	PRN PRN PRN For 1 Doses PRN 6, For 1 Doses PRN PRN	S S S	Duration Until discor Until discor Until discor Until discor Until discor
	acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 4 hours PRN, mild pain (score 1-3), Starting S diphenhydrAMINE (BENADRYL) tablet 25 mg 25 mg, oral, every 6 hours PRN, itching, Starting S diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for minor infusion reaction, Starting S, diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S epINEPHrine (ADRENALIN) 1 mg/1 mL injection 1 mg 1 mg, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg	PRN PRN PRN For 1 Doses PRN 6, For 1 Doses PRN PRN	S S S	