

## Provider Communication

	Interval	Defer Until	Duration
<input type="checkbox"/> Provider communication Prior to beginning Rituxan infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: {Push F2:11554001} drawn on ***.	Every visit	S	Until discount'd

## Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> Sedimentation rate STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd
<input type="checkbox"/> C-reactive protein STAT, Status: Future, Expires: S+366, Clinic Collect	Every visit	S	Until discount'd

## Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discount'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol At start of infusion: obtain vital signs every 15 minutes x 4, then every 30 minutes.	Every visit	S	Until discount'd
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for any signs of infusion reaction. For minor infusion reaction (fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea) stop or slow infusion rate by 50%. If symptoms do not resolve within 5-10 minutes give Benadryl as ordered. Notify MD if symptoms do not resolve within 5-10 minutes. With MD approval, give Solu-medrol as ordered. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor patient closely until completion.	Every visit	S	Until discount'd
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of Severe Infusion Reaction: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension. For severe infusion reaction take the following steps; STOP Infusion and start Normal Saline solution, Notify physician and administer Benadryl as ordered, Monitor vital signs, Administer Epinephrine as ordered. Consult MD for further orders.	Every visit	S	Until discount'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Every visit	S	Until discount'd
<input type="checkbox"/> HEParin, porcine (PF) injection 500 Units 500 Units, intra-catheter, once PRN, line care, Starting S, For 1 Doses Flush portacath before removing needle from portacath.	Every visit	S	Until discount'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discount'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discount'd

## Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discount'd
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discount'd
<input type="checkbox"/> famotidine (PEPCID) tablet 40 mg 40 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discount'd
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg	Every visit	S	Until discount'd

## Pre-Medications (continued)

	Interval	Defer Until	Duration
180 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.			
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 125 mg 125 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd

## Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> RiTUXimab (PF) (RITUXAN) 1,000 mg in sodium chloride 0.9% 250 mL IVPB 1,000 mg, intravenous, once, Starting S, For 1 Doses Do not shake. Start an infusion rate of 50 MG/hour; if there is no infusion reaction, increase the rate by 50 MG/hour increments every 30 minutes, to a maximum rate of 400 MG/hour.	Once	S	1 treatment
<input type="checkbox"/> RiTUXimab (PF) (RITUXAN) 1,000 mg in sodium chloride 0.9% 250 mL IVPB 1,000 mg, intravenous, once, Starting S, For 1 Doses Do not shake. Start an infusion rate of 50 MG/hour; if there is no infusion reaction, increase the rate by 50 MG/hour increments every 30 minutes, to a maximum rate of 400 MG/hour.	Once	S+14	1 treatment
<input type="checkbox"/> RiTUXimab (PF) (RITUXAN) 1,000 mg in sodium chloride 0.9% 1,000 mL INITIAL INFUSION RATE IVPB 1,000 mg, intravenous, once, Starting S, For 1 Doses Initiate infusion at rate of 50 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), then increase infusion rate by 50 mg/hour every 30 minutes, to a maximum rate of 400 mg/hour.	Once	S	1 treatment
<input type="checkbox"/> RiTUXimab (PF) (RITUXAN) 1,000 mg in sodium chloride 0.9% 1,000 mL INITIAL INFUSION RATE IVPB 1,000 mg, intravenous, once, Starting S, For 1 Doses Initiate infusion at rate of 50 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), then increase infusion rate by 50 mg/hour every 30 minutes, to a maximum rate of 400 mg/hour.	Once	S+14	1 treatment

## Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours PRN, mild pain (score 1-3), headaches, fever, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for mild to moderate infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFran) injection 4 mg 4 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discont'd