

## Provider Communication

	Interval	Defer Until	Duration
<input type="checkbox"/> Provider communication Prior to beginning Rituxan infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: {Push F2:11554001} drawn on ***.	Every visit	S	Until discont'd

## Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Every visit	S	Until discont'd
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol Obtain vital signs, prior to Rituximab infusion, then every 15 minutes for 1 hour, then every 30 minutes for 1 hour, then every hour until infusion complete.	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Every visit	S	Until discont'd
<input type="checkbox"/> Discontinue IV Details	Every visit	S	Until discont'd

## Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> famotidine (PEPCID) tablet 40 mg 40 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg 180 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 125 mg 125 mg, intravenous, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Every visit	S	Until discont'd

## Hydration

	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, continuous, Starting S	Every visit	S	Until discont'd

## Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> RiTUXimab (PF) (RITUXAN) 500 mg in sodium chloride 0.9% 500 mL INITIAL INFUSION RATE IVPB 500 mg, intravenous, once, Starting H+30 Minutes, For 1 Doses Initiate infusion at rate of 50 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), then increase infusion rate by 50 mg/hour every 30 minutes, to a maximum rate of 400 mg/hour.	Every visit	S	Until discont'd

## Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, PRN infusion reaction, Starting S, For 1 Doses	PRN	S	Until discont'd

	Interval	Defer Until	Duration
<input type="checkbox"/> epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mg, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinu'd
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, intravenous, once PRN, PRN infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinu'd
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinu'd
<input type="checkbox"/> meperidine (DEMEROL) injection 25 mg 25 mg, intravenous, once PRN, PRN chills and rigors, Starting S, For 1 Doses SLOW IV Push Formulary approved non-pain management indication(s) :	PRN	S	Until discontinu'd