

Pre-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC hemogram Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Prothrombin time with INR Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Partial thromboplastin time, activated Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Fibrinogen Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Ionized calcium Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> LDH Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Hemoglobin electrophoresis Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Miscellaneous referral test Once, Starting S For 1 Occurrences ADAMTS-13 - Misc referral test	Every visit	S	Until discont'd
<input type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1 %) injection 0.3 mL 0.3 mL, injection, once PRN, local anesthetic, Starting S, For 2 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection 0.3 mL 0.3 mL, injection, once PRN, local anesthetic, Starting S, For 2 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> ondansetron ODT (ZOFran-ODT) disintegrating tablet 4 mg 4 mg, oral, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if patient is able to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFran) injection 4 mg 4 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9% 20 mL syringe 12.5 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFran) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 12.5 mg 12.5 mg, oral, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFran) is ineffective and patient is able to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) suppository 12.5 mg 12.5 mg, rectal, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFran) is ineffective and patient is UNable to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 325 mg 325 mg, oral, once PRN, fever, Starting S, For 2 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 0.5 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 1 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

Pre-Procedure (continued)

	Interval	Defer Until	Duration
<input type="checkbox"/> alteplase (CATHFLO) for NON Quinton and Hemodialysis Catheters 1 mg 1 mg, intra-catheter, once, Starting S, For 1 Doses 1 mg dose of Cathflo is for NON Quinton and Hemodialysis Catheter Clearance	Every visit	S	Until discont'd

Procedures

	Interval	Defer Until	Duration
<input type="checkbox"/> LDL apheresis STAT, Once, Starting S For 1 Occurrences Access:	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 5 % (HYPER TONIC) for LDL Apheresis extracorporeal, once, Starting S, For 1 Doses Final Concentration = 5% HYPER TONIC Saline. NOT FOR IV USE!!	Every visit	S	Until discont'd
<input type="checkbox"/> lactated ringer's infusion 1,000 mL 1,000 mL, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> HEParin (porcine) injection 30,000 Units 30,000 Units, extracorporeal, once, Starting S, For 1 Doses Draw up into syringe. Given extracorporeal.	Every visit	S	Until discont'd

Intra-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> ondansetron ODT (ZOFRAN-ODT) disintegrating tablet 4 mg 4 mg, oral, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if patient is able to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFRAN) injection 4 mg 4 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9% 20 mL syringe 12.5 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) tablet 12.5 mg 12.5 mg, oral, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFRAN) is ineffective and patient is able to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) suppository 12.5 mg 12.5 mg, rectal, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 325 mg 325 mg, oral, once PRN, fever, Starting S, For 2 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 0.5 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 1 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

Post-Procedure

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC hemogram Once, Starting S For 1 Occurrences	Every visit	S	Until discont'd
<input type="checkbox"/> Hemoglobin electrophoresis	Every visit	S	Until discont'd

Post-Procedure (continued)

	Interval	Defer Until	Duration
Once, Starting S For 1 Occurrences			
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> ondansetron ODT (ZOFran-ODT) disintegrating tablet 4 mg 4 mg, oral, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if patient is able to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> ondansetron (ZOFran) injection 4 mg 4 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9% 20 mL syringe 12.5 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFran) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.	Every visit	S	Until discont'd
<input type="checkbox"/> promethazine (PHENERGAN) suppository 12.5 mg 12.5 mg, rectal, once PRN, nausea, vomiting, Starting S, For 1 Doses Give if ondansetron (ZOFran) is ineffective and patient is UNable to tolerate oral medication.	Every visit	S	Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 325 mg 325 mg, oral, once PRN, fever, Starting S, For 2 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 0.5 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 1 g, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, once PRN, line care, Starting S, For 1 Doses	Every visit	S	Until discont'd
<input type="checkbox"/> HEParin, porcine (PF) 100 unit/mL injection 100 Units 100 Units, intra-catheter, once PRN, line care, Starting S, For 1 Doses	Every visit	S	Until discont'd

PRN Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> ondansetron (ZOFran) tablet 4 mg 4 mg, oral, once, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses	PRN	S	Until discont'd
<input type="checkbox"/> calcium gluconate injection 1 g, intravenous, once, Starting S, For 1 Doses	PRN	S	Until discont'd