TRANSIENT HEMODIALYSIS THERAPY PLAN

Dialysis Medications

	Interval	Defer Until	Duration
sodium chloride 0.9 % bolus 100 mL 100 mL, intravenous, once, Starting S, For 1 Doses	Every visit	S	Until discont'd

Dialysis Treatment Orders

19313 1				
		Interval	Defer Until	Duration
	Hemodialysis inpatient	Every visit	S	Until discont'o
	Routine, Once, Starting S For 1 Occurrences			
	Duration of Treatment:			
	Fluid removal (L):			
	K+/CA++:			
	Bicarb:			
	Na+:			
	Na+ Modeling:			
	Dialyzer:			
	Dialysate Temperature (C):			
	Blood Flow Rate-As tolerated to a maximum of:			
	Dialysate Flow Rate:			
	Access Site:			
	Hypothermic:			
	Oxygen Therapy - Nasal Cannula; 2 LPM	Every visit	S	Until discont
	Routine, As needed, Starting S For Until specified		-	
	Titrate to keep O2 Sat Above:			
	Device 1: Nasal Cannula			
	Rate in liters per minute: 2 LPM			
	FiO2:			
lysis N	ledications - Heparin (standard)			
		Interval	Defer Until	Duration
	heparin (porcine) injection 1,000-2,000 Units	Every visit	S	Until discont'
	1,000-2,000 Units, intravenous, at 1-2 mL/hr, continuous, Starting S, For 12	Hours		
	heparin (porcine) injection	Every visit	S	Until discont
		,		
	1,000-2,000 Units/hr, intravenous, PRN, Starting S, For 12 Hours			
	1,000-2,000 Units/hr, intravenous, PRN, Starting S, For 12 Hours Initial BOLUS to begin hemodialysis			
lysis N				
lysis N	Initial BOLUS to begin hemodialysis	Interval	Defer Until	Duration
lysis N	Initial BOLUS to begin hemodialysis Iedications - Heparin (mini dose) heparin (porcine) injection 500-1,000 Units	Interval Every visit	Defer Until S	
-	Initial BOLUS to begin hemodialysis Iedications - Heparin (mini dose) heparin (porcine) injection 500-1,000 Units 500-1,000 Units, intravenous, PRN, Starting S, For 12 Hours			
-	Initial BOLUS to begin hemodialysis Iedications - Heparin (mini dose) heparin (porcine) injection 500-1,000 Units			
	Initial BOLUS to begin hemodialysis Iedications - Heparin (mini dose) heparin (porcine) injection 500-1,000 Units 500-1,000 Units, intravenous, PRN, Starting S, For 12 Hours Initial BOLUS to begin hemodialyis	Every visit		Duration Until discont'd Until discont'd
-	Initial BOLUS to begin hemodialysis Iedications - Heparin (mini dose) heparin (porcine) injection 500-1,000 Units 500-1,000 Units, intravenous, PRN, Starting S, For 12 Hours		S	Until discont

Dialysis Medications - Heparin (mini dose) (continued)

Dialysis Medications PRN

	Interval	Defer Until	Duration		
acetaminophen (TYLENOL) tablet 650 mg	PRN	S	Until discont'd		
650 mg, oral, every 4 hours PRN, mild pain, moderate pain, headaches, Starting S, For 12 Hours					
nitroglycerin (NITROSTAT) SL tablet 0.4 mg	PRN	S	Until discont'd		
0.4 mg, sublingual, every 5 min PRN, chest pain, Starting S, For 12 Hours					
May administer up to 3 doses per episode.					